

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use REYATAZ safely and effectively. See full prescribing information for REYATAZ.

REYATAZ[®] (atazanavir) capsules, for oral use

REYATAZ[®] (atazanavir) oral powder

Initial U.S. Approval: 2003

RECENT MAJOR CHANGES

Contraindications (4)

12/2024

INDICATIONS AND USAGE

REYATAZ is a protease inhibitor indicated for use in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and in pediatric patients 3 months and older weighing at least 5 kg. (1)

DOSAGE AND ADMINISTRATION

- **Pretreatment testing:** Renal laboratory testing should be performed in all patients prior to initiation of REYATAZ and continued during treatment with REYATAZ. Hepatic testing should be performed in patients with underlying liver disease prior to initiation of REYATAZ and continued during treatment with REYATAZ. (2.2)
- **Treatment-naïve adults:** REYATAZ 300 mg with ritonavir 100 mg once daily with food or REYATAZ 400 mg once daily with food. (2.3)
- **Treatment-experienced adults:** REYATAZ 300 mg with ritonavir 100 mg once daily with food. (2.3)
- **Pediatric patients:** REYATAZ capsule dosage is based on body weight not to exceed the adult dose and must be taken with food. (2.4)
- **REYATAZ oral powder:** Must be taken with ritonavir and food and should not be used in pediatric patients who weigh less than 5 kg. (2.5)
- **Pregnancy:** REYATAZ 300 mg with ritonavir 100 mg once daily with food, with dosing modifications for some concomitant medications. (2.6)
- **Dosing modifications:** may be required for concomitant therapy (2.3, 2.4, 2.5, 2.6), renal impairment (2.7), and hepatic impairment. (2.8)

DOSAGE FORMS AND STRENGTHS

- Capsules: 200 mg, 300 mg. (3, 16)
- Oral powder: 50 mg packet. (3, 16)

CONTRAINDICATIONS

- In patients with previously demonstrated hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of REYATAZ. (4)
- Coadministration with drugs that are strong inducers of CYP3A, due to the potential for loss of therapeutic effect and development of resistance. (4)
- Coadministration with drugs that are highly dependent on CYP3A or UGT1A1 for clearance, and for which elevated plasma concentrations of the interacting drugs are associated with serious and/or life-threatening events. (4)

WARNINGS AND PRECAUTIONS

- **Cardiac conduction abnormalities:** PR interval prolongation may occur in some patients. ECG monitoring should be considered in patients with preexisting conduction system disease or when administered with other drugs that may prolong the PR interval. (5.1, 7.3, 12.2, 17)

- **Severe Skin Reactions:** Discontinue if severe rash develops. (5.2, 17)
- **Hyperbilirubinemia:** Most patients experience asymptomatic increases in indirect bilirubin, which is reversible upon discontinuation. Do not dose reduce. If a concomitant transaminase increase occurs, evaluate for alternative etiologies. (5.8)
- **Phenylketonuria:** REYATAZ oral powder contains phenylalanine which can be harmful to patients with phenylketonuria. (5.3)
- **Hepatotoxicity:** Patients with hepatitis B or C virus are at risk of increased transaminases or hepatic decompensation. Monitor hepatic laboratory tests prior to therapy and during treatment. (2.8, 5.4, 8.8)
- **Chronic kidney disease** has been reported during postmarketing surveillance in patients with HIV-1 treated with atazanavir, with or without ritonavir. Consider alternatives in patients at high risk for renal disease or with preexisting renal disease. Monitor renal laboratory tests prior to therapy and during treatment. Consider discontinuation of REYATAZ in patients with progressive renal disease. (5.5)
- **Nephrolithiasis and cholelithiasis** have been reported. Consider temporary interruption or discontinuation. (5.6)
- The concomitant use of REYATAZ with ritonavir and certain other medications may result in known or potentially significant drug interactions. Consult the full prescribing information prior to and during treatment for potential drug interactions. (5.7, 7.3)
- Patients receiving REYATAZ may develop new onset or exacerbations of diabetes mellitus/hyperglycemia (5.9), immune reconstitution syndrome (5.10), and redistribution/accumulation of body fat. (5.11)
- **Hemophilia:** Spontaneous bleeding may occur, and additional factor VIII may be required. (5.12)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 2\%$) are nausea, jaundice/scleral icterus, rash, headache, abdominal pain, vomiting, insomnia, peripheral neurologic symptoms, dizziness, myalgia, diarrhea, depression, and fever. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Bristol-Myers Squibb at 1-800-721-5072 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Coadministration of REYATAZ can alter the concentration of other drugs and other drugs may alter the concentration of atazanavir. The potential drug-drug interactions must be considered prior to and during therapy. (4, 7, 12.3)

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** Available human and animal data suggest that atazanavir does not increase the risk of major birth defects overall compared to the background rate. (8.1)
- **Hepatitis B or C co-infection:** Monitor liver enzymes. (5.4, 6.1)
- **Renal impairment:** REYATAZ is not recommended for use in treatment-experienced patients with end-stage renal disease managed with hemodialysis. (2.7, 8.7)
- **Hepatic impairment:** REYATAZ is not recommended in patients with severe hepatic impairment. REYATAZ with ritonavir is not recommended in patients with any degree of hepatic impairment. (2.8, 8.8)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 12/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE	
2	DOSAGE AND ADMINISTRATION	
2.1	Overview	7.1 Potential for REYATAZ to Affect Other Drugs
2.2	Testing Prior to Initiation and During Treatment with REYATAZ	7.2 Potential for Other Drugs to Affect REYATAZ
2.3	Dosage of REYATAZ in Adult Patients	7.3 Established and Other Potentially Significant Drug Interactions
2.4	Dosage of REYATAZ Capsules in Pediatric Patients	7.4 Drugs with No Observed Interactions with REYATAZ
2.5	Dosage and Administration of REYATAZ Oral Powder in Pediatric Patients	8 USE IN SPECIFIC POPULATIONS
2.6	Dosage Adjustments in Pregnant Patients	8.1 Pregnancy
2.7	Dosage in Patients with Renal Impairment	8.2 Lactation
2.8	Dosage Adjustments in Patients with Hepatic Impairment	8.4 Pediatric Use
3	DOSAGE FORMS AND STRENGTHS	8.5 Geriatric Use
4	CONTRAINDICATIONS	8.6 Age/Gender
5	WARNINGS AND PRECAUTIONS	8.7 Impaired Renal Function
5.1	Cardiac Conduction Abnormalities	8.8 Impaired Hepatic Function
5.2	Severe Skin Reactions	10 OVERDOSAGE
5.3	Patients with Phenylketonuria	11 DESCRIPTION
5.4	Hepatotoxicity	12 CLINICAL PHARMACOLOGY
5.5	Chronic Kidney Disease	12.1 Mechanism of Action
5.6	Nephrolithiasis and Cholelithiasis	12.2 Pharmacodynamics
5.7	Risk of Serious Adverse Reactions Due to Drug Interactions	12.3 Pharmacokinetics
5.8	Hyperbilirubinemia	12.4 Microbiology
5.9	Diabetes Mellitus/Hyperglycemia	13 NONCLINICAL TOXICOLOGY
5.10	Immune Reconstitution Syndrome	13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
5.11	Fat Redistribution	14 CLINICAL STUDIES
5.12	Hemophilia	14.1 Adult Participants without Prior Antiretroviral Therapy
5.13	Resistance/Cross-Resistance	14.2 Adult Participants with Prior Antiretroviral Therapy
6	ADVERSE REACTIONS	14.3 Pediatric Participants
6.1	Clinical Trial Experience	16 HOW SUPPLIED/STORAGE AND HANDLING
6.2	Postmarketing Experience	17 PATIENT COUNSELING INFORMATION
7	DRUG INTERACTIONS	

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

REYATAZ[®] is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and in pediatric patients 3 months and older weighing at least 5 kg.

Limitations of Use:

- REYATAZ is not recommended for use in pediatric patients below the age of 3 months due to the risk of kernicterus [see *Use in Specific Populations* (8.4)].
- Use of REYATAZ with ritonavir in treatment-experienced patients should be guided by the number of baseline primary protease inhibitor resistance substitutions [see *Microbiology* (12.4)].

2 DOSAGE AND ADMINISTRATION

2.1 Overview

- REYATAZ capsules and oral powder must be taken with food.
- Do not open the capsules.
- The recommended oral dosage of REYATAZ depends on the treatment history of the patient and the use of other coadministered drugs. When coadministered with H₂-receptor antagonists or proton-pump inhibitors, dose separation may be required [see *Dosage and Administration* (2.3, 2.4, 2.5, and 2.6) and *Drug Interactions* (7)].
- REYATAZ capsules without ritonavir are not recommended for treatment-experienced adult or pediatric patients with prior virologic failure [see *Clinical Studies* (14)].
- REYATAZ oral powder must be taken with ritonavir and is not recommended for use in children who weigh less than 5 kg [see *Dosage and Administration* (2.5)].
- Efficacy and safety of REYATAZ with ritonavir when ritonavir is administered in doses greater than 100 mg once daily have not been established. The use of higher ritonavir doses may alter the safety profile of atazanavir (cardiac effects, hyperbilirubinemia) and, therefore, is not recommended. Prescribers should consult the complete prescribing information for ritonavir when using ritonavir.

2.2 Testing Prior to Initiation and During Treatment with REYATAZ

Renal laboratory testing should be performed in all patients prior to initiation of REYATAZ and continued during treatment with REYATAZ. Renal laboratory testing should include serum creatinine, estimated creatinine clearance, and urinalysis with microscopic examination [see *Warnings and Precautions* (5.5, 5.6)].

Hepatic laboratory testing should be performed in patients with underlying liver disease prior to initiation of REYATAZ and continued during treatment with REYATAZ [see *Warnings and Precautions* (5.4)].

2.3 Dosage of REYATAZ in Adult Patients

Table 1 displays the recommended dosage of REYATAZ capsules in treatment-naïve and treatment-experienced adults. Table 1 also displays recommended dosage of REYATAZ and

ritonavir when given concomitantly with other antiretroviral drugs and H₂-receptor antagonists (H2RA). Ritonavir is required with several REYATAZ dosage regimens (see the ritonavir complete prescribing information about the safe and effective use of ritonavir). The use of REYATAZ in treatment-experienced adult patients without ritonavir is not recommended.

Table 1: Recommended REYATAZ and Ritonavir Dosage in Adults^{a,b}

	REYATAZ Once Daily Dosage	Ritonavir Once Daily Dosage
Treatment-Naive Adult Patients		
recommended regimen	300 mg	100 mg
unable to tolerate ritonavir	400 mg	N/A
in combination with efavirenz	400 mg	100 mg
Treatment-Experienced Adult Patients		
recommended regimen	300 mg	100 mg
in combination with both H2RA and tenofovir DF	400 mg	100 mg

^a See *Drug Interactions* (7) for instructions concerning coadministration of acid-reducing medications (eg, H2RA or proton pump inhibitors [PPIs]), and other antiretroviral drugs (eg, efavirenz, tenofovir DF, and didanosine).

^b For adult patients who cannot swallow the capsules, REYATAZ oral powder is taken once daily with food at the same recommended adult dosage as the capsules along with ritonavir.

2.4 Dosage of REYATAZ Capsules in Pediatric Patients

The recommended daily dosage of REYATAZ capsules and ritonavir in pediatric patients (6 years of age to less than 18 years of age) is based on body weight (see Table 2).

Table 2: Recommended Dosage of REYATAZ Capsules and Ritonavir in Pediatric Patients (6 to less than 18 years of age)^{a,b}

Body weight	REYATAZ Daily Dosage	Ritonavir Daily Dosage
Treatment-Naive and Treatment-Experienced^c		
Less than 15 kg	Capsules not recommended	N/A
At least 15 kg to less than 35 kg	200 mg	100 mg
At least 35 kg	300 mg	100 mg
Treatment-Naive, at least 13 years old and cannot tolerate ritonavir		
At least 40 kg	400 mg	N/A

^a Administer REYATAZ capsules and ritonavir simultaneously with food.

^b The same recommendations regarding the timing and maximum doses of concomitant PPIs and H2RAs in adults also apply to pediatric patients. See *Drug Interactions* (7) for instructions concerning coadministration of acid-

reducing medications (eg, H2RA or PPIs), and other antiretroviral drugs (eg, efavirenz, tenofovir DF, and didanosine).

^c In treatment-experienced patients, REYATAZ capsules must be administered with ritonavir.

When transitioning between formulations, a change in dose may be needed. Consult the dosing table for the specific formulation.

2.5 Dosage and Administration of REYATAZ Oral Powder in Pediatric Patients

REYATAZ oral powder is for use in treatment-naïve or treatment-experienced pediatric patients who are at least 3 months of age and weighing at least 5 kg. REYATAZ oral powder must be mixed with food or a beverage for administration and ritonavir must be given immediately afterwards. Table 3 displays the recommended dosage of REYATAZ oral powder and ritonavir.

Table 3: Recommended Dosage of REYATAZ Oral Powder and Ritonavir in Pediatric Patients (at least 3 months of age and weighing at least 5 kg)^{a,b}

Body Weight	Daily Dosage of REYATAZ Oral Powder	Daily Dosage of Ritonavir Oral Solution
5 kg to less than 15 kg	200 mg (4 packets) ^{c,d}	80 mg
15 kg to less than 25 kg	250 mg (5 packets) ^d	80 mg

^a The same recommendations regarding the timing and maximum doses of concomitant PPIs and H2RAs in adults also apply to pediatric patients. See *Drug Interactions* (7) for instructions concerning coadministration of acid-reducing medications (eg, H2RA or PPIs), and other antiretroviral drugs (eg, efavirenz, tenofovir DF, and didanosine).

^b For pediatric patients at least 25 kg who cannot swallow REYATAZ capsules, 300 mg (6 packets) REYATAZ oral powder is taken once daily with food along with 100 mg ritonavir.

^c Only patients weighing 5 to less than 10 kg who do not tolerate the 200 mg (4 packets) dose of REYATAZ oral powder and have not previously taken an HIV protease inhibitor, may take 150 mg (3 packets) REYATAZ oral powder with close HIV viral load monitoring.

^d Each packet contains 50 mg of REYATAZ.

When transitioning between formulations, a change in dose may be needed. Consult the dosing table for the specific formulation.

Instructions for Mixing REYATAZ Oral Powder [see FDA-approved *Instructions for Use*]

- Determine the number of packets (3, 4, 5 or 6 packets) that are needed.
- Prior to mixing, tap the packet to settle the powder.
- It is preferable to mix REYATAZ oral powder with food such as applesauce or yogurt. Mixing REYATAZ oral powder with a beverage (milk, infant formula, or water) may be used for infants who can drink from a cup. For young infants (less than 6 months) who cannot eat solid

food or drink from a cup, REYATAZ oral powder should be mixed with infant formula and given using an oral dosing syringe. Administration of REYATAZ and infant formula using an infant bottle is not recommended because full dose may not be delivered.

- Use a clean pair of scissors to cut each packet along the dotted line.
- **Mixing with food:** Using a spoon, mix the recommended number of REYATAZ oral powder packets with a minimum of one tablespoon of food (such as applesauce or yogurt). Feed the mixture to the infant or young child. Add an additional one tablespoon of food to the small container, mix, and feed the child the residual mixture.
- **Mixing with a beverage such as milk or water in a small drinking cup:** Using a spoon, mix the recommended number of REYATAZ oral powder packets with a minimum of 30 mL of the beverage. Have the child drink the mixture. Add an additional 15 mL more of beverage to the drinking cup, mix, and have the child drink the residual mixture. If water is used, food should also be taken at the same time.
- **Mixing with liquid infant formula using an oral dosing syringe and a small medicine cup:** Using a spoon, mix the recommended number of REYATAZ oral powder packets with 10 mL of prepared liquid infant formula. Draw up the full amount of the mixture into an oral syringe and administer into either right or left inner cheek of infant. Pour another 10 mL of formula into the medicine cup to rinse off remaining REYATAZ oral powder in cup. Draw up residual mixture into the syringe and administer into either right or left inner cheek of infant.
- Administer ritonavir immediately following REYATAZ powder administration.
- Administer the entire dosage of REYATAZ oral powder (mixed in the food or beverage) within one hour of preparation [may leave the mixture at a temperature of 68°F to 86°F (20°C to 30°C) for up to one hour]. Ensure that the patient eats or drinks all the food or beverage that contains the powder. Additional food may be given after consumption of the entire mixture.

2.6 Dosage Adjustments in Pregnant Patients

Table 4 includes the recommended dosage of REYATAZ capsules and ritonavir in treatment-naïve and treatment-experienced pregnant patients. In these patients, REYATAZ must be administered with ritonavir. There are no dosage adjustments for postpartum patients (see Table 1 for the recommended REYATAZ dosage in adults) [see *Use in Specific Populations* (8.1)].

Table 4: Recommended Dosage of REYATAZ and Ritonavir in Pregnant Patients^a

	REYATAZ Once Daily Dosage	Ritonavir Once Daily Dosage
Treatment-Naïve and Treatment-Experienced		
Recommended Regimen	300 mg	100 mg
Treatment-Experienced During the Second or Third Trimester When Coadministered with either H2RA or Tenofovir DF^b		
In combination with <u>EITHER</u> H2RA <u>OR</u> tenofovir DF	400 mg	100 mg

- ^a See *Drug Interactions (7)* for instructions concerning coadministration of acid-reducing medications (eg, H2RA or PPIs), and other antiretroviral drugs (eg, efavirenz, tenofovir DF, and didanosine).
- ^b REYATAZ is not recommended for treatment-experienced pregnant patients during the second and third trimester taking REYATAZ with **BOTH** tenofovir DF and H2RA.

2.7 Dosage in Patients with Renal Impairment

For patients with renal impairment, including those with severe renal impairment who are not managed with hemodialysis, no dose adjustment is required for REYATAZ. Treatment-naïve patients with end-stage renal disease managed with hemodialysis should receive REYATAZ 300 mg with ritonavir 100 mg. REYATAZ is not recommended in treatment-experienced patients with HIV-1 who have end-stage renal disease managed with hemodialysis [see Use in Specific Populations (8.7)].

2.8 Dosage Adjustments in Patients with Hepatic Impairment

Table 5 displays the recommended REYATAZ dosage in treatment-naïve patients with hepatic impairment. The use of REYATAZ in patients with severe hepatic impairment (Child-Pugh Class C) is not recommended. The coadministration of REYATAZ with ritonavir in patients with any degree of hepatic impairment is not recommended.

Table 5: Recommended Dosage of REYATAZ Capsules in Treatment-Naïve Adults with Hepatic Impairment

	REYATAZ Once Daily Dosage
Mild hepatic impairment (Child-Pugh Class A)	400 mg
Moderate hepatic impairment (Child-Pugh Class B)	300 mg
Severe hepatic impairment (Child-Pugh Class C)	REYATAZ with or without ritonavir is not recommended

3 DOSAGE FORMS AND STRENGTHS

REYATAZ Capsules:

- 200 mg capsule with blue cap and blue body, printed with white ink “BMS 200 mg” on the cap and with white ink “3631” on the body.
- 300 mg capsule with red cap and blue body, printed with white ink “BMS 300 mg” on the cap and with white ink “3622” on the body.

REYATAZ Oral Powder:

- 50 mg of atazanavir as an oral powder in a packet.

4 CONTRAINDICATIONS

REYATAZ is contraindicated:

- in patients with previously demonstrated clinically significant hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of REYATAZ capsules or REYATAZ oral powder [see *Warnings and Precautions* (5.2)].
- when coadministered with drugs that are highly dependent on CYP3A or UGT1A1 for clearance, and for which elevated plasma concentrations of the interacting drugs are associated with serious and/or life-threatening events (see Table 6).
- when coadministered with drugs that are strong inducers of CYP3A due to the potential for loss of therapeutic effect and development of resistance.

Coadministration is contraindicated with, but not limited to, the following drugs listed in Table 6:

Table 6: Drugs Contraindicated with REYATAZ (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Drug Class	Drugs within class that are contraindicated with REYATAZ
Alpha 1-adrenoreceptor antagonist	Alfuzosin
Antiarrhythmics	Amiodarone (with ritonavir), quinidine (with ritonavir)
Anticonvulsants	Carbamazepine, phenobarbital, phenytoin
Antimycobacterials	Rifampin
Antineoplastics	Apalutamide, encorafenib, irinotecan, ivosidenib
Antipsychotics	Lurasidone (with ritonavir), pimozide
Benzodiazepines	Orally administered midazolam ^a , triazolam
Ergot Derivatives	Dihydroergotamine, ergonovine, ergotamine, methylegonovine
GI Motility Agent	Cisapride
Hepatitis C Direct-Acting Antivirals	Elbasvir/grazoprevir; glecaprevir/pibrentasvir
Herbal Products	St. John's wort (<i>Hypericum perforatum</i>)
Lipid-Modifying Agents:	Lomitapide, lovastatin, simvastatin
Phosphodiesterase-5 (PDE-5) Inhibitor	Sildenafil ^b when dosed as REVATIO [®] for the treatment of pulmonary arterial hypertension
Protease Inhibitors	Indinavir
Non-nucleoside Reverse Transcriptase Inhibitors	Nevirapine

^a See *Drug Interactions, Table 16 (7)* for parenterally administered midazolam.

^b See *Drug Interactions, Table 16 (7)* for sildenafil when dosed as VIAGRA[®] for erectile dysfunction.

5 WARNINGS AND PRECAUTIONS

5.1 Cardiac Conduction Abnormalities

REYATAZ has been shown to prolong the PR interval of the electrocardiogram in some study participants. In healthy participants and in participants with HIV-1 treated with atazanavir, abnormalities in atrioventricular (AV) conduction were asymptomatic and generally limited to first-degree AV block. There have been reports of second-degree AV block and other conduction abnormalities [see *Adverse Reactions* (6.2) and *Overdosage* (10)]. In clinical trials that included electrocardiograms, asymptomatic first-degree AV block was observed in 5.9% of atazanavir-treated participants (n=920), 5.2% of lopinavir/ritonavir-treated participants (n=252), 10.4% of nelfinavir-treated participants (n=48), and 3.0% of efavirenz-treated participants (n=329). In Study AI424-045, asymptomatic first-degree AV block was observed in 5% (6/118) of atazanavir with ritonavir-treated participants and 5% (6/116) of lopinavir/ritonavir-treated participants who had on-study electrocardiogram measurements. Because of limited clinical experience in those with preexisting conduction system disease (eg, marked first-degree AV block or second- or third-degree AV block), ECG monitoring should be considered in these patients [see *Clinical Pharmacology* (12.2)].

5.2 Severe Skin Reactions

In controlled clinical trials, rash (all grades, regardless of causality) occurred in approximately 20% of participants with HIV-1 treated with REYATAZ. The median time to onset of rash in clinical studies was 7.3 weeks and the median duration of rash was 1.4 weeks. Rashes were generally mild-to-moderate maculopapular skin eruptions. Treatment-emergent adverse reactions of moderate or severe rash (occurring at a rate of $\geq 2\%$) are presented for the individual clinical studies [see *Adverse Reactions* (6.1)]. Dosing with REYATAZ was often continued without interruption in patients who developed rash. The discontinuation rate for rash in clinical trials was $<1\%$. Cases of Stevens-Johnson syndrome, erythema multiforme, and toxic skin eruptions, including drug rash, eosinophilia, and systemic symptoms (DRESS) syndrome, have been reported in patients receiving REYATAZ [see *Contraindications* (4) and *Adverse Reactions* (6.1)]. REYATAZ should be discontinued if severe rash develops.

5.3 Patients with Phenylketonuria

Phenylalanine can be harmful to patients with phenylketonuria (PKU). REYATAZ oral powder contains phenylalanine (a component of aspartame). Each packet of REYATAZ oral powder contains 35 mg of phenylalanine. REYATAZ capsules do not contain phenylalanine.

5.4 Hepatotoxicity

Patients with underlying hepatitis B or C virus or marked elevations in transaminases before treatment may be at increased risk for developing further transaminase elevations or hepatic decompensation. In these patients, hepatic laboratory testing should be conducted prior to initiating therapy with REYATAZ and during treatment [see *Dosage and Administration* (2.2), *Adverse Reactions* (6.1), and *Use in Specific Populations* (8.8)].

5.5 Chronic Kidney Disease

Chronic kidney disease in patients with HIV-1 treated with atazanavir, with or without ritonavir, has been reported during postmarketing surveillance. Reports included biopsy-proven cases of granulomatous interstitial nephritis associated with the deposition of atazanavir drug crystals in the renal parenchyma. Consider alternatives to REYATAZ in patients at high risk for renal disease or with preexisting renal disease. Renal laboratory testing (including serum creatinine, estimated creatinine clearance, and urinalysis with microscopic examination) should be conducted in all patients prior to initiating therapy with REYATAZ and continued during treatment with REYATAZ. Expert consultation is advised for patients who have confirmed renal laboratory abnormalities while taking REYATAZ. In patients with progressive kidney disease, discontinuation of REYATAZ may be considered [*see Dosage and Administration (2.2 and 2.7) and Adverse Reactions (6.2)*].

5.6 Nephrolithiasis and Cholelithiasis

Cases of nephrolithiasis and/or cholelithiasis have been reported during postmarketing surveillance in patients with HIV-1 receiving REYATAZ therapy. Some patients required hospitalization for additional management, and some had complications. Because these events were reported voluntarily during clinical practice, estimates of frequency cannot be made. If signs or symptoms of nephrolithiasis and/or cholelithiasis occur, temporary interruption or discontinuation of therapy may be considered [*see Adverse Reactions (6.2)*].

5.7 Risk of Serious Adverse Reactions Due to Drug Interactions

Initiation of REYATAZ with ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving REYATAZ with ritonavir, may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of REYATAZ with ritonavir, respectively. These interactions may lead to:

- clinically significant adverse reactions potentially leading to severe, life-threatening, or fatal events from greater exposures of concomitant medications.
- clinically significant adverse reactions from greater exposures of REYATAZ with ritonavir.
- loss of therapeutic effect (virologic response) of REYATAZ with ritonavir and possible development of resistance.

See Table 16 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations [*see Drug Interactions (7)*]. Consider the potential for drug interactions prior to and during therapy containing REYATAZ with ritonavir; and monitor for the adverse reactions associated with concomitant medications [*see Contraindications (4) and Drug Interactions (7)*].

5.8 Hyperbilirubinemia

Most patients taking REYATAZ experience asymptomatic elevations in indirect (unconjugated) bilirubin related to inhibition of UDP-glucuronosyl transferase (UGT). This hyperbilirubinemia is

reversible upon discontinuation of REYATAZ. Hepatic transaminase elevations that occur with hyperbilirubinemia should be evaluated for alternative etiologies. No long-term safety data are available for patients experiencing persistent elevations in total bilirubin >5 times the upper limit of normal (ULN). Alternative antiretroviral therapy to REYATAZ may be considered if jaundice or scleral icterus associated with bilirubin elevations presents cosmetic concerns for patients. Dose reduction of atazanavir is not recommended since long-term efficacy of reduced doses has not been established [see *Adverse Reactions* (6.1)].

5.9 Diabetes Mellitus/Hyperglycemia

New-onset diabetes mellitus, exacerbation of preexisting diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in patients with HIV-1 receiving protease inhibitor therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued protease inhibitor therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and a causal relationship between protease inhibitor therapy and these events has not been established [see *Adverse Reactions* (6.2)].

5.10 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including REYATAZ. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium*, cytomegalovirus, *Pneumocystis jirovecii* pneumonia, or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, Guillain-Barré syndrome, and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.11 Fat Redistribution

Redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and “cushingoid appearance” have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

5.12 Hemophilia

There have been reports of increased bleeding, including spontaneous skin hematomas and hemarthrosis, in patients with hemophilia type A and B treated with protease inhibitors. In some patients, additional factor VIII was given. In more than half of the reported cases, treatment with protease inhibitors was continued or reintroduced. A causal relationship between protease inhibitor therapy and these events has not been established.

5.13 Resistance/Cross-Resistance

Various degrees of cross-resistance among protease inhibitors have been observed. Resistance to atazanavir may not preclude the subsequent use of other protease inhibitors [see *Microbiology (12.4)*].

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- cardiac conduction abnormalities [see *Warnings and Precautions (5.1)*]
- rash [see *Warnings and Precautions (5.2)*]
- hyperbilirubinemia [see *Warnings and Precautions (5.8)*]
- chronic kidney disease [see *Warnings and Precautions (5.5)*]
- nephrolithiasis and cholelithiasis [see *Warnings and Precautions (5.6)*]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions in Treatment-Naive Adult Participants

The safety profile of REYATAZ in treatment-naive adults is based on 1625 participants with HIV-1 in clinical trials. 536 participants received REYATAZ 300 mg with ritonavir 100 mg and 1089 participants received REYATAZ 400 mg or higher (without ritonavir).

The most common adverse reactions were nausea, jaundice/scleral icterus, and rash.

Selected clinical adverse reactions of moderate or severe intensity reported in $\geq 2\%$ of treatment-naive participants receiving combination therapy including REYATAZ 300 mg with ritonavir 100 mg and REYATAZ 400 mg (without ritonavir) are presented in Tables 7 and 8, respectively.

Table 7: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in $\geq 2\%$ of Adult Treatment-Naive Participants with HIV-1,^b Study AI424-138

	96 weeks ^c REYATAZ 300 mg with ritonavir 100 mg (once daily) and tenofovir DF/emtricitabine ^d (n=441)	96 weeks ^c lopinavir/ritonavir ^d 400 mg/ 100 mg (twice daily) and tenofovir DF/emtricitabine ^e (n=437)
Digestive System		
Nausea	4%	8%
Jaundice/scleral icterus	5%	*
Diarrhea	2%	12%

Table 7: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in ≥2% of Adult Treatment-Naive Participants with HIV-1,^b Study AI424-138

Skin and Appendages		
Rash	3%	2%

* None reported in this treatment arm.

^a Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

^b Based on the regimen containing REYATAZ.

^c Median time on therapy.

^d Administered as a fixed-dose.

^e As a fixed-dose product: 300 mg tenofovir DF, 200 mg emtricitabine once daily.

Table 8: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in ≥2% of Adult Treatment-Naive Participants with HIV-1,^b Studies AI424-034, AI424-007, and AI424-008

	Study AI424-034		Studies AI424-007, -008	
	64 weeks^c REYATAZ 400 mg (once daily) with lamivudine/ zidovudine^e (n=404)	64 weeks^c efavirenz 600 mg (once daily) with lamivudine/ zidovudine^e (n=401)	120 weeks^{c,d} REYATAZ 400 mg (once daily) with stavudine and lamivudine or didanosine (n=279)	73 weeks^{c,d} nelfinavir 750 mg TID or 1250 mg BID with stavudine and lamivudine or didanosine (n=191)

Body as a Whole

Headache	6%	6%	1%	2%
----------	----	----	----	----

Digestive System

Nausea	14%	12%	6%	4%
Jaundice/scleral icterus	7%	*	7%	*
Vomiting	4%	7%	3%	3%
Abdominal pain	4%	4%	4%	2%
Diarrhea	1%	2%	3%	16%

Nervous System

Insomnia	3%	3%	<1%	*
Dizziness	2%	7%	<1%	*
Peripheral neurologic symptoms	<1%	1%	4%	3%

Skin and Appendages

Table 8: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in $\geq 2\%$ of Adult Treatment-Naive Participants with HIV-1,^b Studies AI424-034, AI424-007, and AI424-008

	Study AI424-034		Studies AI424-007, -008	
	64 weeks ^c REYATAZ 400 mg (once daily) with lamivudine/ zidovudine ^e (n=404)	64 weeks ^c efavirenz 600 mg (once daily) with lamivudine/ zidovudine ^e (n=401)	120 weeks ^{c,d} REYATAZ 400 mg (once daily) with stavudine and lamivudine or didanosine (n=279)	73 weeks ^{c,d} nelfinavir 750 mg TID or 1250 mg BID with stavudine and lamivudine or didanosine (n=191)
Rash	7%	10%	5%	1%

* None reported in this treatment arm.

^a Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

^b Based on regimens containing REYATAZ.

^c Median time on therapy.

^d Includes long-term follow-up.

^e As a fixed-dose product: 150 mg lamivudine/300 mg zidovudine twice daily.

Adverse Reactions in Treatment-Experienced Adult Participants

The safety profile of REYATAZ in treatment-experienced adults with HIV-1 is based on 119 participants with HIV-1 in clinical trials.

The most common adverse reactions are jaundice/scleral icterus and myalgia.

Selected clinical adverse reactions of moderate or severe intensity reported in $\geq 2\%$ of treatment-experienced participants receiving REYATAZ with ritonavir are presented in Table 9.

Table 9: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in $\geq 2\%$ of Adult Treatment-Experienced Participants with HIV-1,^b Study AI424-045

	48 weeks ^c REYATAZ with ritonavir 300/100 mg (once daily) and tenofovir DF and NRTI (n=119)	48 weeks ^c lopinavir/ritonavir 400/100 mg (twice daily ^d) and tenofovir DF and NRTI (n=118)
Body as a Whole		
Fever	2%	*
Digestive System		

Table 9: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in ≥2% of Adult Treatment-Experienced Participants with HIV-1,^b Study AI424-045

	48 weeks ^c REYATAZ with ritonavir 300/100 mg (once daily) and tenofovir DF and NRTI (n=119)	48 weeks ^c lopinavir/ritonavir 400/100 mg (twice daily ^d) and tenofovir DF and NRTI (n=118)
Jaundice/scleral icterus	9%	*
Diarrhea	3%	11%
Nausea	3%	2%
Nervous System		
Depression	2%	<1%
Musculoskeletal System		
Myalgia	4%	*

* None reported in this treatment arm.

^a Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

^b Based on the regimen containing REYATAZ.

^c Median time on therapy.

^d As a fixed-dose product.

Laboratory Abnormalities in Treatment-Naive Participants

The percentages of adult treatment-naive participants with HIV-1 treated with combination therapy, including REYATAZ 300 mg with ritonavir 100 mg or REYATAZ 400 mg (without ritonavir) with Grade 3–4 laboratory abnormalities, are presented in Tables 10 and 11, respectively.

Table 10: Grade 3–4 Laboratory Abnormalities Reported in ≥2% of Adult Treatment-Naive Participants with HIV-1,^a Study AI424-138

		96 weeks ^b REYATAZ 300 mg with ritonavir 100 mg (once daily) and tenofovir DF/emtricitabine ^c (n=441)	96 weeks ^b lopinavir/ritonavir 400 mg/100 mg ^c (twice daily) and tenofovir DF/emtricitabine ^d (n=437)
Variable	Limit ^e		
Chemistry	<u>High</u>		
SGOT/AST	≥5.1 × ULN	3%	1%

Table 10: Grade 3–4 Laboratory Abnormalities Reported in $\geq 2\%$ of Adult Treatment-Naïve Participants with HIV-1,^a Study AI424-138

Variable	Limit ^e	96 weeks ^b REYATAZ 300 mg with ritonavir 100 mg (once daily) and tenofovir DF/emtricitabine ^c (n=441)	96 weeks ^b lopinavir/ritonavir 400 mg/100 mg ^c (twice daily) and tenofovir DF/emtricitabine ^d (n=437)
SGPT/ALT	$\geq 5.1 \times \text{ULN}$	3%	2%
Total Bilirubin	$\geq 2.6 \times \text{ULN}$	44%	<1%
Lipase	$\geq 2.1 \times \text{ULN}$	2%	2%
Creatine Kinase	$\geq 5.1 \times \text{ULN}$	8%	7%
Total Cholesterol	$\geq 240 \text{ mg/dL}$	11%	25%
Hematology	<u>Low</u>		
Neutrophils	$< 750 \text{ cells/mm}^3$	5%	2%

^a Based on the regimen containing REYATAZ.

^b Median time on therapy.

^c Administered as a fixed-dose product.

^d As a fixed-dose product: 300 mg tenofovir DF, 200 mg emtricitabine once daily.

^e ULN=upper limit of normal.

Table 11: Grade 3–4 Laboratory Abnormalities Reported in $\geq 2\%$ of Adult Treatment-Naïve Participants with HIV-1,^a Studies AI424-034, AI424-007, and AI424-008

Variable	Limit ^d	Study AI424-034		Studies AI424-007, -008	
		64 weeks ^b REYATAZ 400 mg once daily and lamivudine/ zidovudine ^e (n=404)	64 weeks ^b efavirenz 600 mg once daily and lamivudine/ zidovudine ^e (n=401)	120 weeks ^{b,c} REYATAZ 400 mg once daily with stavudine and lamivudine or with stavudine and didanosine (n=279)	73 weeks ^{b,c} nelfinavir 750 mg TID or 1250 mg BID with stavudine and lamivudine or with stavudine and didanosine (n=191)
Chemistry	<u>High</u>				
SGOT/AST	$\geq 5.1 \times \text{ULN}$	2%	2%	7%	5%

Table 11: Grade 3–4 Laboratory Abnormalities Reported in $\geq 2\%$ of Adult Treatment-Naive Participants with HIV-1,^a Studies AI424-034, AI424-007, and AI424-008

Variable	Limit ^d	Study AI424-034		Studies AI424-007, -008	
		64 weeks ^b REYATAZ 400 mg once daily and lamivudine/ zidovudine ^e (n=404)	64 weeks ^b efavirenz 600 mg once daily and lamivudine/ zidovudine ^e (n=401)	120 weeks ^{b,c} REYATAZ 400 mg once daily with stavudine and lamivudine or with stavudine and didanosine (n=279)	73 weeks ^{b,c} nelfinavir 750 mg TID or 1250 mg BID with stavudine and lamivudine or with stavudine and didanosine (n=191)
SGPT/ALT	$\geq 5.1 \times \text{ULN}$	4%	3%	9%	7%
Total Bilirubin	$\geq 2.6 \times \text{ULN}$	35%	<1%	47%	3%
Amylase	$\geq 2.1 \times \text{ULN}$	*	*	14%	10%
Lipase	$\geq 2.1 \times \text{ULN}$	<1%	1%	4%	5%
Creatine Kinase	$\geq 5.1 \times \text{ULN}$	6%	6%	11%	9%
Total Cholesterol	$\geq 240 \text{ mg/dL}$	6%	24%	19%	48%
Triglycerides	$\geq 751 \text{ mg/dL}$	<1%	3%	4%	2%
Hematology	<u>Low</u>				
Hemoglobin	<8.0 g/dL	5%	3%	<1%	4%
Neutrophils	<750 cells/mm ³	7%	9%	3%	7%

* None reported in this treatment arm.

^a Based on regimen(s) containing REYATAZ.

^b Median time on therapy.

^c Includes long-term follow-up.

^d ULN = upper limit of normal.

^e As a fixed-dose product: 150 mg lamivudine, 300 mg zidovudine twice daily.

Change in Lipids from Baseline in Treatment-Naive Participants with HIV-1

For Study AI424-138 and Study AI424-034, changes from baseline in LDL-cholesterol, HDL-cholesterol, total cholesterol, and triglycerides are shown in Tables 12 and 13, respectively.

Table 12: Lipid Values, Mean Change from Baseline, Study AI424-138

	REYATAZ with ritonavir ^{a,b}					lopinavir/ritonavir ^{b,c}				
	Baseline	Week 48		Week 96		Baseline	Week 48		Week 96	
	mg/dL	mg/dL	Change ^d	mg/dL	Change ^d	mg/dL	mg/dL	Change ^d	mg/dL	Change ^d
	(n=428 ^e)	(n=372 ^e)	(n=372 ^e)	(n=342 ^e)	(n=342 ^e)	(n=424 ^e)	(n=335 ^e)	(n=335 ^e)	(n=291 ^e)	(n=291 ^e)
LDL-Cholesterol ^f	92	105	+14%	105	+14%	93	111	+19%	110	+17%
HDL-Cholesterol ^f	37	46	+29%	44	+21%	36	48	+37%	46	+29%
Total Cholesterol ^f	149	169	+13%	169	+13%	150	187	+25%	186	+25%
Triglycerides ^f	126	145	+15%	140	+13%	129	194	+52%	184	+50%

^a REYATAZ 300 mg with ritonavir 100 mg once daily with the fixed-dose product: 300 mg tenofovir DF/ 200 mg emtricitabine once daily.

^b Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 1% in the lopinavir/ritonavir treatment arm and 1% in the REYATAZ with ritonavir arm. Through Week 48, serum lipid-reducing agents were used in 8% in the lopinavir/ritonavir treatment arm and 2% in the REYATAZ with ritonavir arm. Through Week 96, serum lipid-reducing agents were used in 10% in the lopinavir/ritonavir treatment arm and 3% in the REYATAZ with ritonavir arm.

^c Lopinavir/ritonavir (400 mg/100 mg) twice daily with the fixed-dose product 300 mg tenofovir DF/200 mg emtricitabine once daily.

^d The change from baseline is the mean of within-participant changes from baseline for participants with both baseline and Week 48 or Week 96 values and is not a simple difference of the baseline and Week 48 or Week 96 mean values, respectively.

^e Number of participants with LDL-cholesterol measured.

^f Fasting.

Table 13: Lipid Values, Mean Change from Baseline, Study AI424-034

	REYATAZ ^{a,b}			efavirenz ^{b,c}		
	Baseline	Week 48	Week 48	Baseline	Week 48	Week 48
	mg/dL	mg/dL	Change ^d	mg/dL	mg/dL	Change ^d
	(n=383 ^e)	(n=283 ^e)	(n=272 ^e)	(n=378 ^e)	(n=264 ^e)	(n=253 ^e)
LDL-Cholesterol ^f	98	98	+1%	98	114	+18%
HDL-Cholesterol	39	43	+13%	38	46	+24%
Total Cholesterol	164	168	+2%	162	195	+21%
Triglycerides ^f	138	124	-9%	129	168	+23%

^a REYATAZ 400 mg once daily with the fixed-dose product: 150 mg lamivudine, 300 mg zidovudine twice daily.

- ^b Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 0% in the efavirenz treatment arm and <1% in the REYATAZ arm. Through Week 48, serum lipid-reducing agents were used in 3% in the efavirenz treatment arm and 1% in the REYATAZ arm.
- ^c Efavirenz 600 mg once daily with the fixed-dose product: 150 mg lamivudine/300 mg zidovudine twice daily.
- ^d The change from baseline is the mean of within-participant changes from baseline for participants with both baseline and Week 48 values and is not a simple difference of the baseline and Week 48 mean values.
- ^e Number of participants with LDL-cholesterol measured.
- ^f Fasting.

Laboratory Abnormalities in Treatment-Experienced Participants with HIV-1

The percentages of adult treatment-experienced participants with HIV-1 treated with combination therapy, including REYATAZ with ritonavir having Grade 3–4 laboratory abnormalities, are presented in Table 14.

Table 14: Grade 3–4 Laboratory Abnormalities Reported in ≥2% of Adult Treatment-Experienced Participants with HIV-1, Study AI424-045^a

Variable	Limit ^c	48 weeks ^b REYATAZ with ritonavir 300/100 mg (once daily) and tenofovir DF and NRTI (n=119)	48 weeks ^b lopinavir/ritonavir 400/100 mg (twice daily ^d) and tenofovir DF and NRTI (n=118)
Chemistry	<u>High</u>		
SGOT/AST	≥5.1 × ULN	3%	3%
SGPT/ALT	≥5.1 × ULN	4%	3%
Total Bilirubin	≥2.6 × ULN	49%	<1%
Lipase	≥2.1 × ULN	5%	6%
Creatine Kinase	≥5.1 × ULN	8%	8%
Total Cholesterol	≥240 mg/dL	25%	26%
Triglycerides	≥751 mg/dL	8%	12%
Glucose	≥251 mg/dL	5%	<1%
Hematology	<u>Low</u>		
Platelets	<50,000 cells/mm ³	2%	3%
Neutrophils	<750 cells/mm ³	7%	8%

^a Based on regimen(s) containing REYATAZ.

^b Median time on therapy.

^c ULN = upper limit of normal.

^d As a fixed-dose product.

Change in Lipids from Baseline in Treatment-Experienced Participants with HIV-1

For Study AI424-045, changes from baseline in LDL-cholesterol, HDL-cholesterol, total cholesterol, and triglycerides are shown in Table 15. The observed magnitude of dyslipidemia was less with REYATAZ with ritonavir than with lopinavir/ritonavir. However, the clinical impact of such findings has not been demonstrated.

Table 15: Lipid Values, Mean Change from Baseline, Study AI424-045

	REYATAZ with ritonavir ^{a,b}			Lopinavir/ritonavir ^{b,c}		
	Baseline mg/dL (n=111 ^e)	Week 48 mg/dL (n=75 ^e)	Week 48 Change ^d (n=74 ^e)	Baseline mg/dL (n=108 ^e)	Week 48 mg/dL (n=76 ^e)	Week 48 Change ^d (n=73 ^e)
LDL-Cholesterol ^f	108	98	−10%	104	103	+1%
HDL-Cholesterol	40	39	−7%	39	41	+2%
Total Cholesterol	188	170	−8%	181	187	+6%
Triglycerides ^f	215	161	−4%	196	224	+30%

^a REYATAZ 300 mg once daily with ritonavir and tenofovir DF, and 1 NRTI.

^b Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 4% in the lopinavir/ritonavir treatment arm and 4% in the REYATAZ with ritonavir arm. Through Week 48, serum lipid-reducing agents were used in 19% in the lopinavir/ritonavir treatment arm and 8% in the REYATAZ with ritonavir arm.

^c Lopinavir/ritonavir (400/100 mg), as a fixed dose regimen, BID with tenofovir DF and 1 NRTI.

^d The change from baseline is the mean of within-participant changes from baseline for participants with both baseline and Week 48 values and is not a simple difference of the baseline and Week 48 mean values.

^e Number of participants with LDL-cholesterol measured.

^f Fasting.

Adverse Reactions in Pediatric Participants with HIV-1: REYATAZ Capsules

The safety and tolerability of REYATAZ Capsules with and without ritonavir have been established in pediatric participants with HIV-1, at least 6 years of age from the open-label, multicenter clinical trial PACTG 1020A.

The safety profile of REYATAZ in pediatric participants with HIV-1 (6 to less than 18 years of age) taking the capsule formulation was generally similar to that observed in clinical studies of REYATAZ in adults. The most common Grade 2–4 adverse events (≥5%, regardless of causality) reported in pediatric participants were cough (21%), fever (18%), jaundice/scleral icterus (15%), rash (14%), vomiting (12%), diarrhea (9%), headache (8%), peripheral edema (7%), extremity pain (6%), nasal congestion (6%), oropharyngeal pain (6%), wheezing (6%), and rhinorrhea (6%). Asymptomatic second-degree atrioventricular block was reported in <2% of participants. The most

common Grade 3–4 laboratory abnormalities occurring in pediatric participants taking the capsule formulation were elevation of total bilirubin (≥ 3.2 mg/dL, 58%), neutropenia (9%), and hypoglycemia (4%). All other Grade 3–4 laboratory abnormalities occurred with a frequency of less than 3%.

Adverse Reactions in Pediatric Participants with HIV-1: REYATAZ Oral Powder

The data described below reflect exposure to REYATAZ oral powder in 155 participants weighing at least 5 kg to less than 35 kg, including 134 participants exposed for 48 weeks. These data are from two pooled, open-label, multi-center clinical trials in treatment-naïve and treatment-experienced pediatric participants with HIV-1 (AI424-397 [PRINCE I] and AI424-451 [PRINCE II]). Age ranged from 3 months to 10 years of age. In these studies, 51% were female and 49% were male. All participants received ritonavir and 2 nucleoside reverse transcriptase inhibitors (NRTIs).

The safety profile of REYATAZ in pediatric participants taking REYATAZ oral powder was generally similar to that observed in clinical studies of REYATAZ in pediatric participants taking REYATAZ capsules. The most common Grade 3–4 laboratory abnormalities occurring in pediatric participants weighing 5 kg to less than 35 kg taking REYATAZ oral powder were increased amylase (33%), neutropenia (9%), increased SGPT/ALT (9%), elevation of total bilirubin (≥ 2.6 times ULN, 16%), and increased lipase (8%). All other Grade 3–4 laboratory abnormalities occurred with a frequency of less than 3%.

Adverse Reactions in Participants with HIV-1 and Hepatitis B and/or Hepatitis C Virus

In Study AI424-138, 60 participants administered REYATAZ 300 mg with ritonavir 100 mg once daily, and 51 participants treated with lopinavir/ritonavir 400 mg/100 mg (as fixed-dose product) twice daily, each with fixed-dose tenofovir DF/emtricitabine, were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 10% (6/60) of the participants administered REYATAZ with ritonavir and 8% (4/50) of the participants treated with lopinavir/ritonavir. AST levels >5 times ULN developed in 10% (6/60) of the participants administered REYATAZ with ritonavir and none (0/50) of the participants treated with lopinavir/ritonavir.

In Study AI424-045, 20 participants administered REYATAZ 300 mg with ritonavir 100 mg once daily, and 18 participants treated with lopinavir/ritonavir 400 mg/100 mg twice daily (as fixed-dose product), were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 25% (5/20) of the participants administered REYATAZ with ritonavir and 6% (1/18) of the participants treated with lopinavir/ritonavir. AST levels >5 times ULN developed in 10% (2/20) of the participants administered REYATAZ with ritonavir and 6% (1/18) of the participants treated with lopinavir/ritonavir.

In Studies AI424-008 and AI424-034, 74 participants treated with REYATAZ 400 mg once daily, 58 who received efavirenz, and 12 who received nelfinavir were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 15% of the participants treated with REYATAZ, 14% of the participants treated with efavirenz, and 17% of the participants treated with nelfinavir. AST levels >5 times ULN developed in 9% of the participants treated with

REYATAZ, 5% of the participants treated with efavirenz, and 17% of the participants treated with nelfinavir. Within REYATAZ and control regimens, no difference in frequency of bilirubin elevations was noted between seropositive and seronegative participants [see *Warnings and Precautions* (5.8)].

6.2 Postmarketing Experience

The following events have been identified during postmarketing use of REYATAZ. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: edema

Cardiovascular System: second-degree AV block, third-degree AV block, left bundle branch block, QTc prolongation [see *Warnings and Precautions* (5.1)]

Gastrointestinal System: pancreatitis

Hepatic System: hepatic function abnormalities

Hepatobiliary Disorders: cholelithiasis [see *Warnings and Precautions* (5.6)], cholecystitis, cholestasis

Metabolic System and Nutrition Disorders: diabetes mellitus, hyperglycemia [see *Warnings and Precautions* (5.9)]

Musculoskeletal System: arthralgia

Renal System: nephrolithiasis [see *Warnings and Precautions* (5.6)], interstitial nephritis, granulomatous interstitial nephritis, chronic kidney disease [see *Warnings and Precautions* (5.5)]

Skin and Appendages: alopecia, maculopapular rash [see *Contraindications* (4) and *Warnings and Precautions* (5.2)], pruritus, angioedema

7 DRUG INTERACTIONS

7.1 Potential for REYATAZ to Affect Other Drugs

Atazanavir is an inhibitor of CYP3A and UGT1A1. Coadministration of REYATAZ and drugs primarily metabolized by CYP3A or UGT1A1 may result in increased plasma concentrations of the other drug that could increase or prolong its therapeutic and adverse effects.

Atazanavir is a weak inhibitor of CYP2C8. Use of REYATAZ without ritonavir is not recommended when coadministered with drugs highly dependent on CYP2C8 with narrow therapeutic indices (eg, paclitaxel, repaglinide). When REYATAZ with ritonavir is coadministered with substrates of CYP2C8, clinically significant interactions are not expected [see *Clinical Pharmacology*, Table 22 (12.3)].

The magnitude of CYP3A-mediated drug interactions on coadministered drug may change when REYATAZ is coadministered with ritonavir. See the complete prescribing information for ritonavir for information on drug interactions with ritonavir.

7.2 Potential for Other Drugs to Affect REYATAZ

Atazanavir is a CYP3A4 substrate; therefore, drugs that induce CYP3A4 may decrease atazanavir plasma concentrations and reduce REYATAZ's therapeutic effect (*see Table 16*).

Atazanavir solubility decreases as pH increases. Reduced plasma concentrations of atazanavir are expected if proton-pump inhibitors, antacids, buffered medications, or H₂-receptor antagonists are administered with REYATAZ [*see Dosage and Administration (2.3, 2.4, 2.5 and 2.6)*].

7.3 Established and Other Potentially Significant Drug Interactions

Table 16 provides dosing recommendations in adults as a result of drug interactions with REYATAZ. These recommendations are based on either drug interaction studies or predicted interactions due to the expected magnitude of interaction and potential for serious events or loss of efficacy.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>HIV Antiviral Agents</i>		
<i>Nucleoside Reverse Transcriptase Inhibitors (NRTIs):</i> didanosine buffered formulations enteric coated (EC) capsules	↓ atazanavir ↓ didanosine	It is recommended that REYATAZ be given (with food) 2 h before or 1 h after didanosine buffered formulations. Simultaneous administration of didanosine EC and REYATAZ with food results in a decrease in didanosine exposure. Thus, REYATAZ and didanosine EC should be administered at different times.
<i>Nucleotide Reverse Transcriptase Inhibitors:</i> tenofovir disoproxil fumarate (DF)	↓ atazanavir ↑ tenofovir	When coadministered with tenofovir DF in adults, it is recommended that REYATAZ 300 mg be given with ritonavir 100 mg and tenofovir DF 300 mg (all as a single daily dose with food). The mechanism of this interaction is unknown. Higher tenofovir concentrations could potentiate tenofovir-associated adverse reactions, including renal disorders. Patients receiving REYATAZ and tenofovir DF should be monitored for tenofovir-associated adverse reactions. For pregnant patients taking REYATAZ with ritonavir <i>and</i> tenofovir DF, <i>see Dosage and Administration (2.6)</i> .
<i>Non-nucleoside Reverse Transcriptase Inhibitors (NNRTIs):</i> efavirenz	↓ atazanavir	<i>In HIV-treatment-naïve adult patients:</i> If REYATAZ is combined with efavirenz, REYATAZ 400 mg (two 200-mg capsules) should be administered with ritonavir 100 mg simultaneously once daily with food, and efavirenz 600 mg should be administered once daily on an empty stomach, preferably at bedtime. <i>In HIV-treatment-experienced adult patients:</i> Coadministration of REYATAZ with efavirenz is not recommended.
nevirapine	↓ atazanavir ↑ nevirapine	Coadministration of REYATAZ with nevirapine is contraindicated due to the potential loss of virologic response and development of resistance, as well as the potential risk for nevirapine-associated adverse reactions [<i>see Contraindications (4)</i>].

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>Protease Inhibitors:</i> saquinavir (soft gelatin capsules)	↑ saquinavir	Appropriate dosing recommendations for this combination, with or without ritonavir, with respect to efficacy and safety have not been established. In a clinical study, saquinavir 1200 mg coadministered with REYATAZ 400 mg and tenofovir DF 300 mg (all given once daily), and nucleoside analogue reverse transcriptase inhibitors did not provide adequate efficacy [see <i>Clinical Studies</i> (14.2)].
indinavir		Coadministration of REYATAZ with indinavir is contraindicated. Both REYATAZ and indinavir are associated with indirect (unconjugated) hyperbilirubinemia [see <i>Contraindications</i> (4)].
ritonavir	↑ atazanavir	If REYATAZ is coadministered with ritonavir, it is recommended that REYATAZ 300 mg once daily be given with ritonavir 100 mg once daily with food in adults. See the complete prescribing information for ritonavir for information on drug interactions with ritonavir.
Others	↑ other protease inhibitor	Coadministration with other protease inhibitors is not recommended.
<i>Hepatitis C Antiviral Agents</i>		
elbasvir/grazoprevir	↑ grazoprevir	Coadministration of REYATAZ with grazoprevir is contraindicated due to the potential for increased risk of ALT elevations [see <i>Contraindications</i> (4)].
glecaprevir/pibrentasvir	↑ glecaprevir ↑ pibrentasvir	Coadministration of REYATAZ with glecaprevir/pibrentasvir is contraindicated due to the potential for increased the risk of ALT elevations [see <i>Contraindications</i> (4)].
voxilaprevir/sofosbuvir/ velpatasvir	↑ voxilaprevir	Coadministration with REYATAZ is not recommended.
<i>Other Agents</i>		
<i>Alpha 1-Adrenoreceptor Antagonist:</i> alfuzosin	↑ alfuzosin	Coadministration of REYATAZ with alfuzosin is contraindicated due to risk for hypotension [see <i>Contraindications</i> (4)].
<i>Antacids and buffered medications:</i>	↓ atazanavir	REYATAZ should be administered 2 hours before or 1 hour after antacids and buffered medications.
<i>Antiarrhythmics:</i> amiodarone, quinidine amiodarone, bepridil, lidocaine (systemic), quinidine	↑ amiodarone, bepridil, lidocaine (systemic), quinidine	Concomitant use of REYATAZ with ritonavir and either quinidine or amiodarone is contraindicated due to the potential for serious or life-threatening reactions such as cardiac arrhythmias [see <i>Contraindications</i> (4)]. Coadministration with REYATAZ without ritonavir has the potential to produce serious and/or life-threatening adverse events but has not been studied. Caution is warranted and therapeutic concentration monitoring of these drugs is recommended if they are used concomitantly with REYATAZ without ritonavir.
<i>Anticoagulants:</i> warfarin	↑ warfarin	Coadministration with REYATAZ has the potential to produce serious and/or life-threatening bleeding and has not been studied. It is recommended that International Normalized Ratio (INR) be monitored.
<i>Direct-Acting Oral Anticoagulants:</i> betrixaban, dabigatran, edoxaban	↑ betrixaban ↑ dabigatran ↑ edoxaban	Concomitant use of REYATAZ with ritonavir, a strong CYP3A4/P-gp inhibitor, may result in an increased risk of bleeding. Refer to the respective DOAC prescribing information regarding dosing instructions for coadministration with P-gp inhibitors.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
rivaroxaban	REYATAZ with ritonavir ↑ rivaroxaban	Coadministration of REYATAZ with ritonavir, a strong CYP3A4/P-gp inhibitor, and rivaroxaban is not recommended, as it may result in an increased risk of bleeding.
	REYATAZ ↑ rivaroxaban	Coadministration of REYATAZ, a CYP3A4 inhibitor, and rivaroxaban may result in an increased risk of bleeding. Close monitoring is recommended when REYATAZ is coadministered with rivaroxaban.
apixaban	REYATAZ with ritonavir ↑ apixaban	Concomitant use of REYATAZ with ritonavir, a strong CYP3A4/P-gp inhibitor, may result in an increased risk of bleeding. Refer to apixaban dosing instructions for coadministration with strong CYP3A4 and P-gp inhibitors in the apixaban prescribing information.
	REYATAZ ↑ apixaban	Concomitant use of REYATAZ, a CYP3A4 inhibitor, and apixaban may result in an increased risk of bleeding. Close monitoring is recommended when apixaban is coadministered with REYATAZ.
<i>Antidepressants:</i> tricyclic antidepressants	↑ tricyclic antidepressants	Coadministration with REYATAZ has the potential to produce serious and/or life-threatening adverse events and has not been studied. Concentration monitoring of these drugs is recommended if they are used concomitantly with REYATAZ.
trazodone	↑ trazodone	Nausea, dizziness, hypotension, and syncope have been observed following coadministration of trazodone with ritonavir. If trazodone is used with a CYP3A4 inhibitor such as REYATAZ, the combination should be used with caution and a lower dose of trazodone should be considered.
<i>Antiepileptics:</i> carbamazepine	↓ atazanavir ↑ carbamazepine	Coadministration of REYATAZ (with or without ritonavir) with carbamazepine is contraindicated due to the risk for loss of virologic response and development of resistance [see Contraindications (4)].
phenytoin, phenobarbital	↓ atazanavir ↓ phenytoin ↓ phenobarbital	Coadministration of REYATAZ (with or without ritonavir) with phenytoin or phenobarbital is contraindicated due to the risk for loss of virologic response and development of resistance [see Contraindications (4)].
lamotrigine	↓ lamotrigine	Coadministration of lamotrigine and REYATAZ with ritonavir may require dosage adjustment of lamotrigine. No dose adjustment of lamotrigine is required when coadministered with REYATAZ without ritonavir.
<i>Antifungals:</i> ketoconazole, itraconazole	REYATAZ with ritonavir: ↑ ketoconazole ↑ itraconazole	Coadministration of ketoconazole has only been studied with REYATAZ without ritonavir (negligible increase in atazanavir AUC and C _{max}). Due to the effect of ritonavir on ketoconazole, high doses of ketoconazole and itraconazole (>200 mg/day) should be used cautiously when administering REYATAZ with ritonavir.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

<i>Concomitant Drug Class: Specific Drugs</i>	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
voriconazole	REYATAZ with ritonavir in participants with a functional CYP2C19 allele: ↓ voriconazole ↓ atazanavir REYATAZ with ritonavir in participants without a functional CYP2C19 allele: ↑ voriconazole ↓ atazanavir	The use of voriconazole in patients receiving REYATAZ with ritonavir is not recommended unless an assessment of the benefit/risk to the patient justifies the use of voriconazole. Patients should be carefully monitored for voriconazole-associated adverse reactions and loss of either voriconazole or atazanavir efficacy during the coadministration of voriconazole and REYATAZ with ritonavir. Coadministration of voriconazole with REYATAZ (without ritonavir) may affect atazanavir concentrations; however, no data are available.
<i>Antigout:</i> colchicine	↑ colchicine	The coadministration of REYATAZ with colchicine in patients with renal or hepatic impairment is not recommended. Recommended adult dosage of colchicine when administered with REYATAZ: Treatment of gout flares: 0.6 mg (1 tablet) for 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Not to be repeated before 3 days. Prophylaxis of gout flares: If the original regimen was 0.6 mg <i>twice</i> a day, the regimen should be adjusted to 0.3 mg <i>once</i> a day. If the original regimen was 0.6 mg <i>once</i> a day, the regimen should be adjusted to 0.3 mg <i>once every other day</i>. Treatment of familial Mediterranean fever (FMF): Maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day).
<i>Antimycobacterials:</i> rifampin	↓ atazanavir	Coadministration of REYATAZ with rifampin is contraindicated due to the risk for loss of virologic response and development of resistance [see <i>Contraindications (4)</i>].
rifabutin	↑ rifabutin	A rifabutin dose reduction of up to 75% (eg, 150 mg every other day or 3 times per week) is recommended. Increased monitoring for rifabutin-associated adverse reactions including neutropenia is warranted.
<i>Antineoplastics:</i> irinotecan	↑ irinotecan	Coadministration of REYATAZ with irinotecan is contraindicated. Atazanavir inhibits UGT1A1 and may interfere with the metabolism of irinotecan, resulting in increased irinotecan toxicities [see <i>Contraindications (4)</i>].

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
apalutamide	↓ atazanavir	Coadministration of REYATAZ (with or without ritonavir) and apalutamide is contraindicated due to the potential for subsequent loss of virologic response and possible resistance to the class of protease inhibitors [see <i>Contraindications (4)</i>].
ivosidenib	↓ atazanavir ↑ ivosidenib	Coadministration of ivosidenib with REYATAZ (with or without ritonavir) is contraindicated due to the potential for loss of virologic response and risk of serious adverse events such as QT interval prolongation.
encorafenib	↓ atazanavir ↑ encorafenib	Coadministration of encorafenib with REYATAZ (with or without ritonavir) is contraindicated due to the potential for the loss of virologic response and risk of serious adverse events such as QT interval prolongation.
<i>Antiplatelets</i>		
ticagrelor	↑ ticagrelor	Coadministration with ticagrelor is not recommended due to potential increase in the risk of dyspnea, bleeding and other adverse events associated with ticagrelor.
clopidogrel	↓ clopidogrel active metabolite	Coadministration of REYATAZ (with or without ritonavir) and clopidogrel is not recommended. This is due to the potential reduction of the antiplatelet activity of clopidogrel.
<i>Antipsychotics:</i> pimozide	↑ pimozide	Coadministration of REYATAZ with pimozide is contraindicated. This is due to the potential for serious and/or life-threatening reactions such as cardiac arrhythmias [see <i>Contraindications (4)</i>].
lurasidone	REYATAZ with ritonavir ↑ lurasidone REYATAZ ↑ lurasidone	REYATAZ with ritonavir Coadministration of lurasidone with REYATAZ with ritonavir is contraindicated. This is due to the potential for serious and/or life-threatening reactions [see <i>Contraindications (4)</i>]. REYATAZ without ritonavir If coadministration is necessary, reduce the lurasidone dose. Refer to the lurasidone prescribing information for concomitant use with moderate CYP3A4 inhibitors.
quetiapine	↑ quetiapine	Initiation of REYATAZ with ritonavir in patients taking quetiapine: Consider alternative antiretroviral therapy to avoid increases in quetiapine exposures. If coadministration is necessary, reduce the quetiapine dose to 1/6 of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for recommendations on adverse reaction monitoring. Initiation of quetiapine in patients taking REYATAZ with ritonavir: Refer to the quetiapine prescribing information for initial dosing and titration of quetiapine.
<i>Benzodiazepines:</i> midazolam (oral) triazolam	↑ midazolam ↑ triazolam	Coadministration of REYATAZ with either orally administered midazolam or triazolam is contraindicated. Triazolam and orally administered midazolam are extensively metabolized by CYP3A4, and coadministration with REYATAZ can lead to the potential for serious and/or life-threatening events such as prolonged or increased sedation or respiratory depression [see <i>Contraindications (4)</i>].

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
parenterally administered midazolam ^b	↑ midazolam	Coadministration with parenteral midazolam should be done in a setting which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dosage reduction for midazolam should be considered, especially if more than a single dose of midazolam is administered.
<i>Calcium channel blockers:</i> diltiazem	↑ diltiazem and desacetyl-diltiazem	Caution is warranted. A dose reduction of diltiazem by 50% should be considered. ECG monitoring is recommended. Coadministration of diltiazem and REYATAZ with ritonavir has not been studied.
felodipine, nifedipine, nicardipine, and verapamil	↑ calcium channel blocker	Caution is warranted. Dose titration of the calcium channel blocker should be considered. ECG monitoring is recommended.
<i>Corticosteroids:</i> dexamethasone and other corticosteroids (<i>all routes of administration</i>)	↓ atazanavir ↑ corticosteroids	Coadministration with dexamethasone or other corticosteroids that induce CYP3A may result in loss of therapeutic effect of REYATAZ and development of resistance to atazanavir and/or ritonavir. Alternative corticosteroids should be considered. Coadministration with corticosteroids (all routes of administration) that are metabolized by CYP3A, particularly for long-term use, may increase the risk for development of systemic corticosteroid effects including Cushing's syndrome and adrenal suppression. Consider the potential benefit of treatment versus the risk of systemic corticosteroid effects. For coadministration of cutaneously administered corticosteroids sensitive to CYP3A inhibition, refer to the prescribing information of the corticosteroid for additional information.
<i>Endothelin receptor antagonists:</i> bosentan	REYATAZ ↓ atazanavir REYATAZ with ritonavir ↑ bosentan	Coadministration of bosentan and REYATAZ without ritonavir is not recommended. For adult patients who have been receiving REYATAZ with ritonavir for at least 10 days, start bosentan at 62.5 mg once daily or every other day based on individual tolerability. For adult patients who have been receiving bosentan, discontinue bosentan at least 36 hours before starting REYATAZ with ritonavir. At least 10 days after starting REYATAZ with ritonavir, resume bosentan at 62.5 mg once daily or every other day based on individual tolerability.
<i>Ergot derivatives:</i> dihydroergotamine, ergotamine, ergonovine, methylexgonovine	↑ ergot derivatives	Coadministration of REYATAZ with ergot derivatives is contraindicated. This is due to the potential for serious and/or life-threatening events such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues [see <i>Contraindications (4)</i>].
<i>GI Motility Agents:</i> cisapride	↑ cisapride	Coadministration of REYATAZ with cisapride is contraindicated. This is due to the potential for serious and/or life-threatening reactions such as cardiac arrhythmias [see <i>Contraindications (4)</i>].
<i>Gonadotropin-releasing hormone Receptor (GnRH) Antagonists:</i> elagolix	↓ atazanavir ↑ elagolix	Coadministration of elagolix and REYATAZ with or without ritonavir is not recommended due to the potential of loss of virologic response and the potential risk of adverse events such as bone loss and hepatic transaminase elevations associated with elagolix. In the event coadministration is necessary, limit concomitant use of elagolix 200mg twice daily with REYATAZ with or without ritonavir for up to 1 month or limit concomitant use of elagolix 150 mg once daily with REYATAZ (with or without ritonavir) for up to 6 months and monitor virologic response.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>Herbal Products:</i> St. John's wort (<i>Hypericum perforatum</i>)	↓ atazanavir	Coadministration of products containing St. John's wort with REYATAZ is contraindicated. This may result in loss of therapeutic effect of REYATAZ and the development of resistance [see <i>Contraindications (4)</i>].
<i>Kinase inhibitors:</i> fostamatinib	↑ R406 (active metabolite of fostamatinib)	When coadministering fostamatinib with REYATAZ (with or without ritonavir), monitor for toxicities of R406 exposure resulting in dose-related adverse events such as hepatotoxicity and neutropenia. Fostamatinib dose reduction may be required.
<i>Lipid-modifying agents</i> <i>HMG-CoA reductase inhibitors:</i> lovastatin, simvastatin	↑ lovastatin ↑ simvastatin	Coadministration of REYATAZ with lovastatin or simvastatin is contraindicated. This is due to the potential for serious reactions such as myopathy, including rhabdomyolysis [see <i>Contraindications (4)</i>].
atorvastatin, rosuvastatin	↑ atorvastatin ↑ rosuvastatin	Titrate atorvastatin dose carefully and use the lowest necessary dose. Rosuvastatin dose should not exceed 10 mg/day. The risk of myopathy, including rhabdomyolysis, may be increased when HIV protease inhibitors, including REYATAZ, are used in combination with these drugs.
<i>Other Lipid Modifying Agents:</i> lomitapide	↑ lomitapide	Coadministration of REYATAZ with lomitapide is contraindicated. This is due to the potential for risk of markedly increased transaminase levels and hepatotoxicity associated with increased plasma concentrations of lomitapide. The mechanism of interaction is CYP3A4 inhibition by atazanavir and/or ritonavir [see <i>Contraindications (4)</i>].

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>H₂-Receptor antagonists</i>	↓ atazanavir	Coadministration may result in loss of virologic response and development of resistance. <i>In HIV-treatment-naïve adult patients:</i> REYATAZ 300 mg with ritonavir 100 mg once daily with food should be administered simultaneously with, and/or at least 10 hours after, a dose of the H₂-receptor antagonist (H2RA). An H2RA dose comparable to famotidine 20 mg once daily up to a dose comparable to famotidine 40 mg twice daily can be used with REYATAZ 300 mg with ritonavir 100 mg in treatment-naïve patients. OR For patients unable to tolerate ritonavir, REYATAZ 400 mg once daily with food should be administered at least 2 hours before and at least 10 hours after a dose of the H2RA. No single dose of the H2RA should exceed a dose comparable to famotidine 20 mg, and the total daily dose should not exceed a dose comparable to famotidine 40 mg. The use of REYATAZ without ritonavir in pregnant patients is not recommended. <i>In treatment-experienced adult patients:</i> Whenever an H2RA is given to a patient receiving REYATAZ with ritonavir, the H2RA dose should not exceed a dose comparable to famotidine 20 mg twice daily, and the REYATAZ with ritonavir doses should be administered simultaneously with, and/or at least 10 hours after, the dose of the H2RA. • REYATAZ 300 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with an H2RA. • REYATAZ 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with both tenofovir DF and an H2RA. • REYATAZ 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with either tenofovir DF or an H2RA for pregnant patients during the second and third trimester. REYATAZ is not recommended for pregnant patients during the second and third trimester taking REYATAZ with both tenofovir DF and an H2RA.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>Hormonal contraceptives:</i> ethinyl estradiol and norgestimate or norethindrone	↓ ethinyl estradiol ↑ norgestimate^c ↑ ethinyl estradiol ↑ norethindrone^d	Use caution if considering coadministration of oral contraceptives with REYATAZ or REYATAZ with ritonavir. If REYATAZ with ritonavir is coadministered with an oral contraceptive, it is recommended that the oral contraceptive contain at least 35 mcg of ethinyl estradiol. If REYATAZ is administered without ritonavir, the oral contraceptive should contain no more than 30 mcg of ethinyl estradiol. Potential safety risks include substantial increases in progesterone exposure. The long-term effects of increases in concentration of the progestational agent are unknown and could increase the risk of insulin resistance, dyslipidemia, and acne. Coadministration of REYATAZ or REYATAZ with ritonavir and other hormonal contraceptives (eg, contraceptive patch, contraceptive vaginal ring, or injectable contraceptives) or oral contraceptives containing progestogens other than norethindrone or norgestimate, or less than 25 mcg of ethinyl estradiol, has not been studied; therefore, alternative methods of contraception are recommended.
<i>Immunosuppressants:</i> cyclosporine, sirolimus, tacrolimus	↑ immunosuppressants	Therapeutic concentration monitoring is recommended for these immunosuppressants when coadministered with REYATAZ.
<i>Inhaled beta agonist:</i> salmeterol	↑ salmeterol	Coadministration of salmeterol with REYATAZ is not recommended. Concomitant use of salmeterol and REYATAZ may result in increased risk of cardiovascular adverse reactions associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia.
<i>Inhaled/nasal steroid:</i> fluticasone	REYATAZ ↑ fluticasone REYATAZ with ritonavir ↑ fluticasone	Concomitant use of fluticasone propionate and REYATAZ without ritonavir should be used with caution. Consider alternatives to fluticasone propionate, particularly for long-term use. With concomitant use of fluticasone propionate and REYATAZ with ritonavir systemic corticosteroid effects, including Cushing's syndrome and adrenal suppression, have been reported during postmarketing use in patients receiving ritonavir and inhaled or intranasally administered fluticasone propionate. Coadministration of fluticasone propionate and REYATAZ with ritonavir is not recommended unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects [see <i>Warnings and Precautions</i> (5.1)].
<i>Macrolide antibiotics:</i> clarithromycin	↑ clarithromycin ↓ 14-OH clarithromycin ↑ atazanavir	Increased concentrations of clarithromycin may cause QTc prolongations; therefore, a dose reduction of clarithromycin by 50% should be considered when it is coadministered with REYATAZ. In addition, concentrations of the active metabolite 14-OH clarithromycin are significantly reduced; consider alternative therapy for indications other than infections due to <i>Mycobacterium avium</i> complex. Coadministration of REYATAZ with ritonavir and clarithromycin has not been studied.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>Opioids:</i> buprenorphine	REYATAZ or REYATAZ with ritonavir ↑ buprenorphine ↑ norbuprenorphine REYATAZ ↓ atazanavir	Coadministration of REYATAZ with ritonavir and buprenorphine warrants clinical monitoring for sedation and cognitive effects. A dose reduction of buprenorphine may be considered. The coadministration of REYATAZ and buprenorphine without ritonavir is not recommended.
<i>PDE5 inhibitors:</i> sildenafil, tadalafil, vardenafil	↑ sildenafil ↑ tadalafil ↑ vardenafil	Coadministration with REYATAZ has not been studied but may result in an increase in PDE5 inhibitor-associated adverse reactions, including hypotension, syncope, visual disturbances, and priapism. Use of PDE5 inhibitors for pulmonary arterial hypertension (PAH): Coadministration of REYATAZ with REVATIO[®] (sildenafil) for the treatment of pulmonary hypertension (PAH) is contraindicated [see <i>Contraindications (4)</i>]. The following dose adjustments are recommended for the use of ADCIRCA[®] (tadalafil) with REYATAZ: Coadministration of ADCIRCA[®] in patients on REYATAZ (with or without ritonavir): - For patients receiving REYATAZ (with or without ritonavir) for at least one week, start ADCIRCA[®] at 20 mg once daily. Increase to 40 mg once daily based on individual tolerability. Coadministration of REYATAZ (with or without ritonavir) in patients on ADCIRCA[®]: - Avoid the use of ADCIRCA[®] when starting REYATAZ (with or without ritonavir). Stop ADCIRCA[®] at least 24 hours before starting REYATAZ (with or without ritonavir). At least one week after starting REYATAZ (with or without ritonavir), resume ADCIRCA[®] at 20 mg once daily. Increase to 40 mg once daily based on individual tolerability. Use of PDE5 inhibitors for erectile dysfunction: Use VIAGRA[®] (sildenafil) with caution at reduced doses of 25 mg every 48 hours with increased monitoring for adverse events. Use CIALIS[®] (tadalafil) with caution at reduced doses of 10 mg every 72 hours with increased monitoring for adverse events. REYATAZ with ritonavir: Use vardenafil with caution at reduced doses of no more than 2.5 mg every 72 hours with increased monitoring for adverse reactions. REYATAZ: Use vardenafil with caution at reduced doses of no more than 2.5 mg every 24 hours with increased monitoring for adverse reactions.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies^a or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>Proton-pump inhibitors: omeprazole</i>	↓ atazanavir	Coadministration of REYATAZ with or without ritonavir and omeprazole may result in loss of virologic response and development of resistance. <i>In HIV-treatment-naïve adult patients:</i> The proton-pump inhibitor (PPI) dose should not exceed a dose comparable to omeprazole 20 mg and must be taken approximately 12 hours prior to the REYATAZ 300 mg with ritonavir 100 mg dose. <i>In HIV-treatment-experienced adult patients:</i> Coadministration of REYATAZ with PPIs is not recommended.

^a For magnitude of interactions see *Clinical Pharmacology, Tables 21 and 22 (12.3)*.

^b See *Contraindications (4), Table 6* for orally administered midazolam.

^c In combination with atazanavir 300 mg with ritonavir 100 mg once daily.

^d In combination with atazanavir 400 mg once daily.

7.4 Drugs with No Observed Interactions with REYATAZ

No clinically significant drug interactions were observed when REYATAZ was coadministered with methadone, fluconazole, acetaminophen, atenolol, or the nucleoside reverse transcriptase inhibitors lamivudine or zidovudine [see *Clinical Pharmacology, Tables 21 and 22 (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in patients exposed to REYATAZ during pregnancy. Healthcare providers are encouraged to register patients by calling the Antiretroviral Pregnancy Registry (APR) at 1-800-258-4263.

Risk Summary

Atazanavir has been evaluated in a limited number of women during pregnancy. Available human and animal data suggest that atazanavir does not increase the risk of major birth defects overall compared to the background rate [see *Data*]. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively. No treatment-related malformations were observed in rats and rabbits, for which the atazanavir exposures were 0.7-1.2 times of those at the human clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir). When atazanavir was administered to

rats during pregnancy and throughout lactation, reversible neonatal growth retardation was observed [see Data].

Clinical Considerations

Dose Adjustments during Pregnancy and the Postpartum Period

- REYATAZ must be administered with ritonavir in pregnant patients.
- For pregnant patients, no dosage adjustment is required for REYATAZ with the following exceptions:
 - For treatment-experienced pregnant women during the second or third trimester, when REYATAZ is coadministered with either an H₂-receptor antagonist **or** tenofovir DF, REYATAZ 400 mg with ritonavir 100 mg once daily is recommended. There are insufficient data to recommend a REYATAZ dose for use with both an H₂-receptor antagonist *and* tenofovir DF in treatment-experienced pregnant patients.
- No dosage adjustment is required for postpartum patients. However, patients should be closely monitored for adverse events because atazanavir exposures could be higher during the first 2 months after delivery [see Dosage and Administration (2.6) and Clinical Pharmacology (12.3)].

Maternal Adverse Reactions

Cases of lactic acidosis syndrome, sometimes fatal, and symptomatic hyperlactatemia have occurred in pregnant women using REYATAZ in combination with nucleoside analogues, which are associated with an increased risk of lactic acidosis syndrome.

Hyperbilirubinemia occurs frequently in patients who take REYATAZ [see Warnings and Precautions (5.8)], including those who are pregnant [see Data].

Advise pregnant women of the potential risks of lactic acidosis syndrome and hyperbilirubinemia.

Fetal/Neonatal Adverse Reactions

All infants, including neonates exposed to REYATAZ *in utero*, should be monitored for the development of severe hyperbilirubinemia during the first few days of life [see Data].

Data

Human Data

In Study AI424-182, REYATAZ with ritonavir (300/100 mg or 400/100 mg) coadministered with lamivudine/zidovudine (150 mg/ 300 mg, as fixed-dose product) was administered to 41 pregnant women with HIV-1, during the second or third trimester. Among the 39 women who completed the study, 38 women achieved an HIV-1 RNA less than 50 copies/mL at time of delivery. Six of 20 (30%) women on REYATAZ with ritonavir 300/100 mg and 13 of 21 (62%) women on REYATAZ with ritonavir 400/100 mg experienced hyperbilirubinemia (total bilirubin greater than or equal to 2.6 times ULN). There were no cases of lactic acidosis observed in clinical trial AI424-182.

Atazanavir drug concentrations in fetal umbilical cord blood were approximately 12% to 19% of maternal concentrations. Among the 40 infants born to 40 pregnant women with HIV-1, all had

test results that were negative for HIV-1 DNA at the time of delivery and/or during the first 6 months postpartum. All 40 infants received antiretroviral prophylactic treatment containing zidovudine. No evidence of severe hyperbilirubinemia (total bilirubin levels greater than 20 mg/dL) or acute or chronic bilirubin encephalopathy was observed among neonates in this study. However, 10/36 (28%) infants (6 greater than or equal to 38 weeks gestation and 4 less than 38 weeks gestation) had bilirubin levels of 4 mg/dL or greater within the first day of life.

Lack of ethnic diversity was a study limitation. In the study population, 33/40 (83%) infants were Black/African American, who have a lower incidence of neonatal hyperbilirubinemia than Caucasians and Asians. In addition, women with Rh incompatibility were excluded, as well as women who had a previous infant who developed hemolytic disease and/or had neonatal pathologic jaundice (requiring phototherapy).

Additionally, of the 38 infants who had glucose samples collected in the first day of life, 3 had adequately collected serum glucose samples with values of less than 40 mg/dL that could not be attributed to maternal glucose intolerance, difficult delivery, or sepsis.

Based on prospective reports from the APR of approximately 1600 live births following exposure to atazanavir-containing regimens (including 1037 live births in infants exposed in the first trimester and 569 exposed in second/third trimesters), there was no difference between atazanavir, and overall birth defects compared with the background birth defect rate. In the U.S. general population, the estimated background risk of major birth defects in clinically recognized pregnancies is 2-4%.

Animal Data

In animal reproduction studies, there was no evidence of mortality or teratogenicity in offspring born to animals at systemic drug exposure levels (AUC) 0.7 (in rabbits) to 1.2 (in rats) times those observed at the human clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir). In pre- and postnatal development studies in the rat, atazanavir caused neonatal growth retardation during lactation that reversed after weaning. Maternal drug exposure at this dose was 1.3 times the human exposure at the recommended clinical exposure. Minimal maternal toxicity occurred at this exposure level.

8.2 Lactation

Risk Summary

Atazanavir has been detected in human milk. No data are available regarding atazanavir effects on milk production. Atazanavir was present in the milk of lactating rats and was associated with neonatal growth retardation that reversed after weaning.

Potential risks of breastfeeding include: (1) HIV-1 transmission (in infants without HIV-1), (2) developing viral resistance (in infants with HIV-1), and (3) adverse reactions in a breastfed infant similar to those seen in adults.

8.4 Pediatric Use

REYATAZ is indicated in combination with other antiretroviral agents for the treatment of pediatric patients with HIV-1, 3 months of age and older weighing at least 5 kg. REYATAZ is not recommended for use in pediatric patients below the age of 3 months due to the risk of kernicterus [see *Indications and Usage (1)*]. All REYATAZ contraindications, warnings, and precautions apply to pediatric patients [see *Contraindications (4)* and *Warnings and Precautions (5)*].

The safety, pharmacokinetic profile, and virologic response of REYATAZ in pediatric patients at least 3 months of age and older weighing at least 5 kg were established in three open-label, multicenter clinical trials: PACTG 1020A, AI424-451, and AI424-397 [see *Clinical Pharmacology (12.3)* and *Clinical Studies (14.3)*]. The safety profile in pediatric patients was generally similar to that observed in adults [see *Adverse Reactions (6.1)*]. See *Dosage and Administration (2.4, 2.5)* for dosing recommendations for the use of REYATAZ capsules and REYATAZ oral powder in pediatric patients.

8.5 Geriatric Use

Clinical studies of REYATAZ did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Based on a comparison of mean single-dose pharmacokinetic values for C_{max} and AUC, a dose adjustment based upon age is not recommended. In general, appropriate caution should be exercised in the administration and monitoring of REYATAZ in elderly patients reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Age/Gender

A study of the pharmacokinetics of atazanavir was performed in young (n=29; 18-40 years) and elderly (n=30; ≥65 years) healthy participants. There were no clinically significant pharmacokinetic differences observed due to age or gender.

8.7 Impaired Renal Function

REYATAZ is not recommended for use in treatment-experienced patients with HIV-1, who have end-stage renal disease managed with hemodialysis [see *Dosage and Administration (2.7)* and *Clinical Pharmacology (12.3)*].

8.8 Impaired Hepatic Function

REYATAZ is not recommended for use in patients with severe hepatic impairment. REYATAZ with ritonavir is not recommended in patients with any degree of hepatic impairment [see *Dosage and Administration (2.8)* and *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

Human experience of acute overdose with REYATAZ is limited. Single doses up to 1200 mg (three times the 400 mg maximum recommended dose) have been taken by healthy participants without symptomatic untoward effects. A single self-administered overdose of 29.2 g of REYATAZ in a patient with HIV-1 (73 times the 400-mg recommended dose) was associated with asymptomatic bifascicular block and PR interval prolongation. These events resolved


spontaneously. At REYATAZ doses resulting in high atazanavir exposures, jaundice due to indirect (unconjugated) hyperbilirubinemia (without associated liver function test changes) or PR interval prolongation may be observed [see *Warnings and Precautions* (5.1, 5.8) and *Clinical Pharmacology* (12.2)].

Treatment of overdosage with REYATAZ should consist of general supportive measures, including monitoring of vital signs and ECG, and observations of the patient's clinical status. If indicated, elimination of unabsorbed atazanavir should be achieved by emesis or gastric lavage. Administration of activated charcoal may also be used to aid removal of unabsorbed drug. There is no specific antidote for overdose with REYATAZ. Since atazanavir is extensively metabolized by the liver and is highly protein bound, dialysis is unlikely to be beneficial in significant removal of this medicine.

11 DESCRIPTION

The active ingredient in REYATAZ capsules and oral powder is atazanavir sulfate, which is an HIV-1 protease inhibitor.

The chemical name for atazanavir sulfate is (3*S*,8*S*,9*S*,12*S*)-3,12-Bis(1,1-dimethylethyl)-8-hydroxy-4,11-dioxo-9-(phenylmethyl)-6-[[4-(2-pyridinyl)phenyl]methyl]-2,5,6,10,13-pentaazatetradecanedioic acid dimethyl ester, sulfate (1:1). Its molecular formula is $C_{38}H_{52}N_6O_7 \cdot H_2SO_4$, which corresponds to a molecular weight of 802.9 (sulfuric acid salt). The free base molecular weight is 704.9. Atazanavir sulfate has the following structural formula:



Atazanavir sulfate is a white to pale-yellow crystalline powder. It is slightly soluble in water (4-5 mg/mL, free base equivalent) with the pH of a saturated solution in water being about 1.9 at $24 \pm 3^\circ C$.

REYATAZ Capsules are available for oral administration in strengths of 200 mg or 300 mg of atazanavir, which are equivalent to 227.8 mg or 341.69 mg of atazanavir sulfate, respectively. The capsules also contain the following inactive ingredients: crospovidone, lactose monohydrate, and magnesium stearate. The capsule shells contain the following inactive ingredients: gelatin, FD&C Blue No. 2, titanium dioxide, black iron oxide, red iron oxide, and yellow iron oxide. The capsules are printed with ink containing shellac, titanium dioxide, FD&C Blue No. 2, isopropyl alcohol, ammonium hydroxide, propylene glycol, n-butyl alcohol, simethicone, and dehydrated alcohol.

REYATAZ oral powder comes in a packet containing 50 mg of atazanavir equivalent to 56.9 mg of atazanavir sulfate in 1.5 g of powder. The powder is off-white to pale yellow and contains the following inactive ingredients: aspartame, sucrose, and orange-vanilla flavor.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Atazanavir is an HIV-1 antiretroviral drug [see *Microbiology (12.4)*].

12.2 Pharmacodynamics

Cardiac Electrophysiology

Concentration- and dose-dependent prolongation of the PR interval in the electrocardiogram has been observed in healthy participants receiving atazanavir. In placebo-controlled Study AI424-076, the mean ($\pm$ SD) maximum change in PR interval from the predose value was 24 ($\pm$ 15) msec following oral dosing with 400 mg of atazanavir (n=65) compared to 13 ($\pm$ 11) msec following dosing with placebo (n=67). The PR interval prolongations in this study were asymptomatic. There is limited information on the potential for a pharmacodynamic interaction in humans between atazanavir and other drugs that prolong the PR interval of the electrocardiogram [see *Warnings and Precautions (5.1)*].

Electrocardiographic effects of atazanavir were determined in a clinical pharmacology study of 72 healthy participants. Oral doses of 400 mg (maximum recommended dosage) and 800 mg (twice the maximum recommended dosage) were compared with placebo; there was no concentration-dependent effect of atazanavir on the QTc interval (using Fridericia's correction). In 1793 participants with HIV-1, receiving antiretroviral regimens, QTc prolongation was comparable in the atazanavir and comparator regimens. No atazanavir-treated healthy participant or participant with HIV-1 in clinical trials had a QTc interval >500 msec [see *Warnings and Precautions (5.1)*].

12.3 Pharmacokinetics

The pharmacokinetics of atazanavir were evaluated in adult participants who either were healthy, or with HIV-1, after administration of REYATAZ 400 mg once daily and after administration of REYATAZ 300 mg with ritonavir 100 mg once daily (see Table 17).

Table 17: Steady-State Pharmacokinetics of Atazanavir in Healthy Participants or Participants with HIV-1 in the Fed State


Parameter	400 mg once daily		300 mg with ritonavir 100 mg once daily	
	Healthy Participants (n=14)	Participants with HIV-1 (n=13)	Healthy Participants (n=28)	Participants with HIV-1 (n=10)
C_{max} (ng/mL)				
Geometric mean (CV%)	5199 (26)	2298 (71)	6129 (31)	4422 (58)
Mean (SD)	5358 (1371)	3152 (2231)	6450 (2031)	5233 (3033)
T_{max} (h)				
Median	2.5	2.0	2.7	3.0
AUC (ng•h/mL)				
Geometric mean (CV%)	28132 (28)	14874 (91)	57039 (37)	46073 (66)
Mean (SD)	29303 (8263)	22262 (20159)	61435 (22911)	53761 (35294)
T-half (h)				
Mean (SD)	7.9 (2.9)	6.5 (2.6)	18.1 (6.2) ^a	8.6 (2.3)
C_{min} (ng/mL)				
Geometric mean (CV%)	159 (88)	120 (109)	1227 (53)	636 (97)
Mean (SD)	218 (191)	273 (298) ^b	1441 (757)	862 (838)

^a n=26.

^b n=12.

Figure 1 displays the mean plasma concentrations of atazanavir at steady state after REYATAZ 400 mg once daily (as two 200-mg capsules) with a light meal and after REYATAZ 300 mg (as two 150-mg capsules) with ritonavir 100 mg once daily with a light meal in adult participants with HIV-1.

Figure 1: Mean (SD) Steady-State Plasma Concentrations of Atazanavir 400 mg (n=13) and 300 mg with Ritonavir (n=10) for Adult Participants with HIV-1



Absorption

Atazanavir is rapidly absorbed with a T_{max} of approximately 2.5 hours. Atazanavir demonstrates nonlinear pharmacokinetics with greater than dose-proportional increases in AUC and C_{max} values over the dose range of 200 to 800 mg once daily. Steady state is achieved between Days 4 and 8, with an accumulation of approximately 2.3-fold.

Food Effect

Administration of REYATAZ with food enhances bioavailability and reduces pharmacokinetic variability. Administration of a single 400-mg dose of REYATAZ with a light meal (357 kcal, 8.2 g fat, 10.6 g protein) resulted in a 70% increase in AUC and 57% increase in C_{max} relative to the fasting state. Administration of a single 400-mg dose of REYATAZ with a high-fat meal (721 kcal, 37.3 g fat, 29.4 g protein) resulted in a mean increase in AUC of 35% with no change in C_{max} relative to the fasting state. Administration of REYATAZ with either a light meal or high-fat meal decreased the coefficient of variation of AUC and C_{max} by approximately one-half compared to the fasting state.

Coadministration of a single 300-mg dose of REYATAZ and a 100-mg dose of ritonavir with a light meal (336 kcal, 5.1 g fat, 9.3 g protein) resulted in a 33% increase in the AUC and a 40% increase in both the C_{max} and the 24-hour concentration of atazanavir relative to the fasting state. Coadministration with a high-fat meal (951 kcal, 54.7 g fat, 35.9 g protein) did not affect the AUC of atazanavir relative to fasting conditions and the C_{max} was within 11% of fasting values. The 24-hour concentration following a high-fat meal was increased by approximately 33% due to delayed absorption; the median T_{max} increased from 2.0 to 5.0 hours. Coadministration of

REYATAZ with ritonavir with either a light or a high-fat meal decreased the coefficient of variation of AUC and $C_{\max}$ by approximately 25% compared to the fasting state.

Distribution

Atazanavir is 86% bound to human serum proteins and protein binding is independent of concentration. Atazanavir binds to both alpha-1-acid glycoprotein (AAG) and albumin to a similar extent (89% and 86%, respectively). In a multiple-dose study in participants with HIV-1 dosed with REYATAZ 400 mg once daily with a light meal for 12 weeks, atazanavir was detected in the cerebrospinal fluid and semen. The cerebrospinal fluid/plasma ratio for atazanavir (n=4) ranged between 0.0021 and 0.0226 and seminal fluid/plasma ratio (n=5) ranged between 0.11 and 4.42.

Metabolism

Atazanavir is extensively metabolized in humans. The major biotransformation pathways of atazanavir in humans consisted of monooxygenation and dioxygenation. Other minor biotransformation pathways for atazanavir or its metabolites consisted of glucuronidation, N-dealkylation, hydrolysis, and oxygenation with dehydrogenation. Two minor metabolites of atazanavir in plasma have been characterized. Neither metabolite demonstrated *in vitro* antiviral activity. *In vitro* studies using human liver microsomes suggested that atazanavir is metabolized by CYP3A.

Elimination

Following a single 400-mg dose of ^{14}C -atazanavir, 79% and 13% of the total radioactivity was recovered in the feces and urine, respectively. Unchanged drug accounted for approximately 20% and 7% of the administered dose in the feces and urine, respectively. The mean elimination half-life of atazanavir in healthy participants (n=214) and adult participants with HIV-1 (n=13) was approximately 7 hours at steady state following a dose of 400 mg daily with a light meal.

Specific Populations

Renal Impairment

In healthy participants, the renal elimination of unchanged atazanavir was approximately 7% of the administered dose. REYATAZ has been studied in adult participants with severe renal impairment (n=20), including those on hemodialysis, at multiple doses of 400 mg once daily. The mean atazanavir $C_{\max}$ was 9% lower, AUC was 19% higher, and $C_{\min}$ was 96% higher in participants with severe renal impairment not undergoing hemodialysis (n=10), than in age-, weight-, and gender-matched participants with normal renal function. In a 4-hour dialysis session, 2.1% of the administered dose was removed. When atazanavir was administered either prior to, or following hemodialysis (n=10), the geometric means for $C_{\max}$, AUC, and $C_{\min}$ were approximately 25% to 43% lower compared to participants with normal renal function. The mechanism of this decrease is unknown. REYATAZ is not recommended for use in treatment-experienced patients with HIV-1 who have end-stage renal disease managed with hemodialysis [see *Dosage and Administration* (2.7)].

Hepatic Impairment

REYATAZ has been studied in adult participants with moderate-to-severe hepatic impairment (14 with Child-Pugh B and 2 with Child-Pugh C) after a single 400-mg dose. The mean $AUC_{(0-\infty)}$ was 42% greater in participants with impaired hepatic function than in healthy participants. The mean half-life of atazanavir in hepatically impaired participants was 12.1 hours compared to 6.4 hours in healthy participants. A dose reduction to 300 mg is recommended for patients with moderate hepatic impairment (Child-Pugh Class B) who have not experienced prior virologic failure as increased concentrations of atazanavir are expected. REYATAZ is not recommended for use in patients with severe hepatic impairment. The pharmacokinetics of REYATAZ in combination with ritonavir has not been studied in participants with hepatic impairment; thus, coadministration of REYATAZ with ritonavir is not recommended for use in patients with any degree of hepatic impairment [see *Dosage and Administration* (2.8)].

Pediatrics

The pharmacokinetic parameters for atazanavir at steady state in pediatric participants taking the powder formulation are summarized in Table 18 by weight ranges [see *Dosage and Administration* (2.5)].

Table 18: Steady-State Pharmacokinetics of Atazanavir (powder formulation) with Ritonavir in Pediatric Participants with HIV-1

Body Weight (range in kg) [n]	atazanavir with ritonavir Dose (mg)	C _{max} ng/mL Geometric Mean (CV%)	AUC ng•h/mL Geometric Mean (CV%)	C _{min} ng/mL Geometric Mean (CV%)
5 to <10 [20]	150/80	4131 (55%)	32503 (61%)	336 (76%)
5 to <10 [10]	200/80	4466 (59%)	39519 (54%)	550 (60%)
10 to <15 [18]	200/80	5197 (53%)	50305 (67%)	572 (111%)
15 to <25 [32]	250/80	5394 (46%)	55687 (45%)	686 (68%)
25 to <35 [8]	300/100	4209 (52%)	44329 (63%)	468 (104%)

The pharmacokinetic parameters for atazanavir at steady state in pediatric participants taking the capsule formulation were predicted by a population pharmacokinetic model and are summarized in Table 19 by weight ranges that correspond to the recommended doses [see *Dosage and Administration* (2.4)].

Table 19: Predicted Steady-State Pharmacokinetics of Atazanavir (capsule formulation) with Ritonavir in Pediatric Participants with HIV-1

Body Weight (range in kg)	atazanavir with ritonavir Dose (mg)	C _{max} ng/mL Geometric Mean (CV%)	AUC ng•h/mL Geometric Mean (CV%)	C _{min} ng/mL Geometric Mean (CV%)
15 to <35	200/100	3303 (86%)	37235 (84%)	538 (99%)
≥35	300/100	2980 (82%)	37643 (83%)	653 (89%)

Pregnancy

The pharmacokinetic data from pregnant women with HIV-1 receiving REYATAZ Capsules with ritonavir are presented in Table 20.

Table 20: Steady-State Pharmacokinetics of Atazanavir with Ritonavir in Pregnant Women with HIV-1 in the Fed State

Pharmacokinetic Parameter	Atazanavir 300 mg with ritonavir 100 mg		
	2nd Trimester (n=5 ^a)	3rd Trimester (n=20)	Postpartum ^b (n=34)
C _{max} ng/mL	3078.85	3291.46	5721.21
Geometric mean (CV%)	(50)	(48)	(31)
AUC ng•h/mL	27657.1	34251.5	61990.4
Geometric mean (CV%)	(43)	(43)	(32)
C _{min} ng/mL ^c	538.70	668.48	1462.59
Geometric mean (CV%)	(46)	(50)	(45)

^a Available data during the 2nd trimester are limited.

^b Atazanavir peak concentrations and AUCs were found to be approximately 28% to 43% higher during the postpartum period (4-12 weeks) than those observed historically in, non-pregnant patients with HIV-1. Atazanavir plasma trough concentrations were approximately 2.2-fold higher during the postpartum period when compared to those observed historically in non-pregnant patients with HIV-1.

^c C_{min} is concentration 24 hours post-dose.

Drug Interaction Data

Atazanavir is a metabolism-dependent CYP3A inhibitor, with a K_{inact} value of 0.05 to 0.06 min⁻¹ and K_i value of 0.84 to 1.0 μM. Atazanavir is also a direct inhibitor for UGT1A1 (K_i=1.9 μM) and CYP2C8 (K_i=2.1 μM).

Atazanavir has been shown *in vivo* not to induce its own metabolism nor to increase the biotransformation of some drugs metabolized by CYP3A. In a multiple-dose study, REYATAZ decreased the urinary ratio of endogenous 6β-OH cortisol to cortisol versus baseline, indicating that CYP3A production was not induced.

Clinically significant interactions are not expected between atazanavir and substrates of CYP2C19, CYP2C9, CYP2D6, CYP2B6, CYP2A6, CYP1A2, or CYP2E1. Clinically significant interactions are not expected between atazanavir when administered with ritonavir and substrates of CYP2C8. See the complete prescribing information for ritonavir for information on other potential drug interactions with ritonavir.

Based on known metabolic profiles, clinically significant drug interactions are not expected between REYATAZ and dapsone, trimethoprim/sulfamethoxazole, azithromycin, or erythromycin. REYATAZ does not interact with substrates of CYP2D6 (eg, nortriptyline, desipramine, metoprolol).

Drug interaction studies were performed with REYATAZ and other drugs likely to be coadministered and some drugs commonly used as probes for pharmacokinetic interactions. The effects of coadministration of REYATAZ on the AUC, C_{max} , and C_{min} are summarized in Tables 21 and 22. Neither didanosine EC nor diltiazem had a significant effect on atazanavir exposures (see Table 22 for effect of atazanavir on didanosine EC or diltiazem exposures). REYATAZ did not have a significant effect on the exposures of didanosine (when administered as the buffered tablet), stavudine, or fluconazole. For information regarding clinical recommendations, see *Drug Interactions (7)*.

Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C_{max}	AUC	C_{min}
atenolol	50 mg QD, d 7–11 (n=19) and d 19–23	400 mg QD, d 1–11 (n=19)	1.00 (0.89, 1.12)	0.93 (0.85, 1.01)	0.74 (0.65, 0.86)
clarithromycin	500 mg BID, d 7–10 (n=29) and d 18–21	400 mg QD, d 1–10 (n=29)	1.06 (0.93, 1.20)	1.28 (1.16, 1.43)	1.91 (1.66, 2.21)
didanosine (ddl) (buffered tablets) and stavudine (d4T) ^b	ddl: 200 mg × 1 dose, d4T: 40 mg × 1 dose (n=31)	400 mg × 1 dose simultaneously with ddl and d4T (n=31)	0.11 (0.06, 0.18)	0.13 (0.08, 0.21)	0.16 (0.10, 0.27)
	ddl: 200 mg × 1 dose, d4T: 40 mg × 1 dose (n=32)	400 mg × 1 dose 1 h after ddl + d4T (n=32)	1.12 (0.67, 1.18)	1.03 (0.64, 1.67)	1.03 (0.61, 1.73)
efavirenz	600 mg QD, d 7–20 (n=27)	400 mg QD, d 1–20 (n=27)	0.41 (0.33, 0.51)	0.26 (0.22, 0.32)	0.07 (0.05, 0.10)
	600 mg QD, d 7–20 (n=13)	400 mg QD, d 1–6 (n=23) then 300 mg with ritonavir 100 mg QD, 2 h before efavirenz, d 7–20 (n=13)	1.14 (0.83, 1.58)	1.39 (1.02, 1.88)	1.48 (1.24, 1.76)

Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C _{max}	AUC	C _{min}
	600 mg QD, d 11–24 (pm) (n=14)	300 mg QD with ritonavir 100 mg QD, d 1–10 (pm) (n=22), then 400 mg QD with ritonavir 100 mg QD, d 11–24 (pm), (simultaneously with efavirenz) (n=14)	1.17 (1.08, 1.27)	1.00 (0.91, 1.10)	0.58 (0.49, 0.69)
famotidine	40 mg BID, d 7–12 (n=15)	400 mg QD, d 1–6 (n=45), d 7–12 (simultaneous administration) (n=15)	0.53 (0.34, 0.82)	0.59 (0.40, 0.87)	0.58 (0.37, 0.89)
	40 mg BID, d 7–12 (n=14)	400 mg QD (pm), d 1–6 (n=14), d 7–12 (10 h after, 2 h before famotidine) (n=14)	1.08 (0.82, 1.41)	0.95 (0.74, 1.21)	0.79 (0.60, 1.04)
	40 mg BID, d 11–20 (n=14) ^c	300 mg QD with ritonavir 100 mg QD, d 1–10 (n=46), d 11–20 ^d (simultaneous administration) (n=14)	0.86 (0.79, 0.94)	0.82 (0.75, 0.89)	0.72 (0.64, 0.81)
	20 mg BID, d 11–17 (n=18)	300 mg QD with ritonavir 100 mg QD and tenofovir DF 300 mg QD, d 1–10 (am) (n=39), d 11–17 (am) (simultaneous administration with am famotidine) (n=18) ^{d,e}	0.91 (0.84, 0.99)	0.90 (0.82, 0.98)	0.81 (0.69, 0.94)
	40 mg QD (pm), d 18–24 (n=20)	300 mg QD with ritonavir 100 mg QD and tenofovir DF 300 mg QD, d 1–10 (am) (n=39), d 18–24 (am) (12 h after pm famotidine) (n=20) ^e	0.89 (0.81, 0.97)	0.88 (0.80, 0.96)	0.77 (0.63, 0.93)
	40 mg BID, d 18–24 (n=18)	300 mg QD with ritonavir 100 mg QD and tenofovir DF 300 mg QD, d 1–10 (am) (n=39), d 18–24 (am) (10 h after pm famotidine and 2 h before am famotidine) (n=18) ^e	0.74 (0.66, 0.84)	0.79 (0.70, 0.88)	0.72 (0.63, 0.83)

Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C _{max}	AUC	C _{min}
	40 mg BID, d 11–20 (n=15)	300 mg QD with ritonavir 100 mg QD, d 1–10 (am) (n=46), then 400 mg QD with ritonavir 100 mg QD, d 11–20 (am) (n=15)	1.02 (0.87, 1.18)	1.03 (0.86, 1.22)	0.86 (0.68, 1.08)
grazoprevir/ elbasvir	grazoprevir 200 mg QD d 1–35 (n = 11)	300 mg QD with ritonavir 100 mg QD, d 1–35 (n = 11)	1.12 (1.01, 1.24)	1.43 (1.30, 1.57)	1.23 (1.13, 1.34)
	elbasvir 50 mg QD d 1–35 (n = 8)	300 mg QD with ritonavir 100 mg QD, d 1–35 (n = 8)	1.02 (0.96, 1.08)	1.07 (0.98, 1.17)	1.15 (1.02, 1.29)
ketoconazole	200 mg QD, d 7–13 (n=14)	400 mg QD, d 1–13 (n=14)	0.99 (0.77, 1.28)	1.10 (0.89, 1.37)	1.03 (0.53, 2.01)
nevirapine ^{f,g}	200 mg BID, d 1–23 (n=23)	300 mg QD with ritonavir 100 mg QD, d 4–13, then 400 mg QD with ritonavir 100 mg QD, d 14–23 (n=23) ^h	0.72 (0.60, 0.86)	0.58 (0.48, 0.71)	0.28 (0.20, 0.40)
			1.02 (0.85, 1.24)	0.81 (0.65, 1.02)	0.41 (0.27, 0.60)
omeprazole	40 mg QD, d 7–12 (n=16) ⁱ	400 mg QD, d 1–6 (n=48), d 7–12 (n=16)	0.04 (0.04, 0.05)	0.06 (0.05, 0.07)	0.05 (0.03, 0.07)
	40 mg QD, d 11–20 (n=15) ⁱ	300 mg QD with ritonavir 100 mg QD, d 1–20 (n=15)	0.28 (0.24, 0.32)	0.24 (0.21, 0.27)	0.22 (0.19, 0.26)
	20 mg QD, d 17–23 (am) (n=13)	300 mg QD with ritonavir 100 mg QD, d 7–16 (pm) (n=27), d 17–23 (pm) (n=13) ^{j,k}	0.61 (0.46, 0.81)	0.58 (0.44, 0.75)	0.54 (0.41, 0.71)
	20 mg QD, d 17–23 (am) (n=14)	300 mg QD with ritonavir 100 mg QD, d 7–16 (am) (n=27), then 400 mg QD with ritonavir 100 mg QD, d 17–23 (am) (n=14) ^{l,m}	0.69 (0.58, 0.83)	0.70 (0.57, 0.86)	0.69 (0.54, 0.88)
pitavastatin	4 mg QD for 5 days	300 mg QD for 5 days	1.13 (0.96, 1.32)	1.06 (0.90, 1.26)	NA
rifabutin	150 mg QD, d 15–28 (n=7)	400 mg QD, d 1–28 (n=7)	1.34 (1.14, 1.59)	1.15 (0.98, 1.34)	1.13 (0.68, 1.87)

Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C _{max}	AUC	C _{min}
rifampin	600 mg QD, d 17–26 (n=16)	300 mg QD with ritonavir 100 mg QD, d 7–16 (n=48), d 17–26 (n=16)	0.47 (0.41, 0.53)	0.28 (0.25, 0.32)	0.02 (0.02, 0.03)
ritonavir ^b	100 mg QD, d 11–20 (n=28)	300 mg QD, d 1–20 (n=28)	1.86 (1.69, 2.05)	3.38 (3.13, 3.63)	11.89 (10.23, 13.82)
tenofovir DF ^c	300 mg QD, d 9–16 (n=34)	400 mg QD, d 2–16 (n=34)	0.79 (0.73, 0.86)	0.75 (0.70, 0.81)	0.60 (0.52, 0.68)
	300 mg QD, d 15–42 (n=10)	300 mg with ritonavir 100 mg QD, d 1–42 (n=10)	0.72 ^p (0.50, 1.05)	0.75 ^p (0.58, 0.97)	0.77 ^p (0.54, 1.10)
voriconazole (Participants with at least one functional CYP2C19 allele)	200 mg BID, d 2–3, 22–30; 400 mg BID, d 1, 21 (n=20)	300 mg with ritonavir 100 mg QD, d 11–30 (n=20)	0.87 (0.80, 0.96)	0.88 (0.82, 0.95)	0.80 (0.72, 0.90)
voriconazole (Participants without a functional CYP2C19 allele)	50 mg BID, d 2–3, 22–30; 100 mg BID, d 1, 21 (n=8)	300 mg with ritonavir 100 mg QD, d 11–30 (n=8)	0.81 (0.66, 1.00)	0.80 (0.65, 0.97)	0.69 (0.54, 0.87)

^a Data provided are under fed conditions unless otherwise noted.

^b All drugs were given under fasted conditions.

^c REYATAZ 300 mg with ritonavir 100 mg once daily coadministered with famotidine 40 mg twice daily resulted in atazanavir geometric mean C_{max} that was similar and AUC and C_{min} values that were 1.79- and 4.46-fold higher relative to REYATAZ 400 mg once daily alone.

^d Similar results were noted when famotidine 20 mg BID was administered 2 hours after and 10 hours before atazanavir 300 mg with ritonavir 100 mg and tenofovir DF 300 mg.

^e Coadministration of atazanavir with ritonavir and tenofovir DF was administered after a light meal.

^f Study was conducted in participants with HIV-1.

^g Compared with atazanavir 400 mg historical data without nevirapine (n=13), the ratio of geometric means (90% confidence intervals) for C_{max}, AUC, and C_{min} were 1.42 (0.98, 2.05), 1.64 (1.11, 2.42), and 1.25 (0.66, 2.36), respectively, for atazanavir with ritonavir 300/100 mg; and 2.02 (1.42, 2.87), 2.28 (1.54, 3.38), and 1.80 (0.94, 3.45), respectively, for atazanavir with ritonavir 400/100 mg.

^h Parallel group design; n=23 for atazanavir with ritonavir and nevirapine, n=22 for atazanavir 300 mg/ritonavir 100 mg without nevirapine. Participants were treated with nevirapine prior to study entry.

ⁱ Omeprazole 40 mg was administered on an empty stomach 2 hours before REYATAZ.

- ^j Omeprazole 20 mg was administered 30 minutes prior to a light meal in the morning and REYATAZ 300 mg with ritonavir 100 mg in the evening after a light meal, separated by 12 hours from omeprazole.
- ^k REYATAZ 300 mg with ritonavir 100 mg once daily separated by 12 hours from omeprazole 20 mg daily resulted in increases in atazanavir geometric mean AUC (10%) and C_{min} (2.4-fold), with a decrease in C_{max} (29%) relative to REYATAZ 400 mg once daily in the absence of omeprazole (study days 1–6).
- ^l Omeprazole 20 mg was given 30 minutes prior to a light meal in the morning and REYATAZ 400 mg with ritonavir 100 mg once daily after a light meal, 1 hour after omeprazole. Effects on atazanavir concentrations were similar when REYATAZ 400 mg with ritonavir 100 mg was separated from omeprazole 20 mg by 12 hours.
- ^m REYATAZ 400 mg with ritonavir 100 mg once daily administered with omeprazole 20 mg once daily resulted in increases in atazanavir geometric mean AUC (32%) and C_{min} (3.3-fold), with a decrease in C_{max} (26%) relative to REYATAZ 400 mg once daily in the absence of omeprazole (study days 1–6).
- ⁿ Compared with atazanavir 400 mg QD historical data, administration of atazanavir with ritonavir 300/100 mg QD increased the atazanavir geometric mean values of C_{max} , AUC, and C_{min} by 18%, 103%, and 671%, respectively.
- ^o Note that similar results were observed in studies where administration of tenofovir DF and REYATAZ was separated by 12 hours.
- ^p Ratio of atazanavir with ritonavir and tenofovir DF to atazanavir with ritonavir. Atazanavir 300 mg with ritonavir 100 mg results in higher atazanavir exposure than atazanavir 400 mg (see footnote ^o). The geometric mean values of atazanavir pharmacokinetic parameters when coadministered with ritonavir and tenofovir DF were: C_{max} = 3190 ng/mL, AUC = 34459 ng•h/mL, and C_{min} = 491 ng/mL. Study was conducted in participants with HIV-1.
- NA = not available.

Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			C_{max}	AUC	C_{min}
acetaminophen	1 g BID, d 1–20 (n=10)	300 mg QD with ritonavir 100 mg QD, d 11–20 (n=10)	0.87 (0.77, 0.99)	0.97 (0.91, 1.03)	1.26 (1.08, 1.46)
atenolol	50 mg QD, d 7–11 (n=19) and d 19–23	400 mg QD, d 1–11 (n=19)	1.34 (1.26, 1.42)	1.25 (1.16, 1.34)	1.02 (0.88, 1.19)
clarithromycin	500 mg BID, d 7–10 (n=21) and d 18–21	400 mg QD, d 1–10 (n=21)	1.50 (1.32, 1.71) OH-clarithromycin: 0.28 (0.24, 0.33)	1.94 (1.75, 2.16) OH-clarithromycin: 0.30 (0.26, 0.34)	2.60 (2.35, 2.88) OH-clarithromycin: 0.38 (0.34, 0.42)
ddI (enteric-coated [EC] capsules) ^b	400 mg d 1 (fasted), d 8 (fed) (n=34)	400 mg QD, d 2–8 (n=34)	0.64 (0.55, 0.74)	0.66 (0.60, 0.74)	1.13 (0.91, 1.41)
	400 mg d 1 (fasted), d 19 (fed) (n=31)	300 mg QD with ritonavir 100 mg QD, d 9–19 (n=31)	0.62 (0.52, 0.74)	0.66 (0.59, 0.73)	1.25 (0.92, 1.69)

Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			C _{max}	AUC	C _{min}
diltiazem	180 mg QD, d 7–11 (n=28) and d 19–23	400 mg QD, d 1–11 (n=28)	1.98 (1.78, 2.19) desacetyl-diltiazem: 2.72 (2.44, 3.03)	2.25 (2.09, 2.16) desacetyl-diltiazem: 2.65 (2.45, 2.87)	2.42 (2.14, 2.73) desacetyl-diltiazem: 2.21 (2.02, 2.42)
ethinyl estradiol & norethindrone ^c	Ortho-Novum [®] 7/7/7 QD, d 1–29 (n=19)	400 mg QD, d 16–29 (n=19)	ethinyl estradiol: 1.15 (0.99, 1.32) norethindrone: 1.67 (1.42, 1.96)	ethinyl estradiol: 1.48 (1.31, 1.68) norethindrone: 2.10 (1.68, 2.62)	ethinyl estradiol: 1.91 (1.57, 2.33) norethindrone: 3.62 (2.57, 5.09)
ethinyl estradiol & norgestimate ^d	Ortho Tri-Cyclen [®] QD, d 1–28 (n=18), then Ortho Tri-Cyclen [®] LO QD, d 29–42 ^e (n=14)	300 mg QD with ritonavir 100 mg QD, d 29–42 (n=14)	ethinyl estradiol: 0.84 (0.74, 0.95) 17-deacetyl norgestimate: ^f 1.68 (1.51, 1.88)	ethinyl estradiol: 0.81 (0.75, 0.87) 17-deacetyl norgestimate: ^f 1.85 (1.67, 2.05)	ethinyl estradiol: 0.63 (0.55, 0.71) 17-deacetyl norgestimate: ^f 2.02 (1.77, 2.31)
glecaprevir/pibrentasvir	300 mg glecaprevir (n=12)	300 mg QD with ritonavir 100 mg QD (n=12)	≥4.06 ^g (3.15, 5.23)	≥6.53 ^g (5.24, 8.14)	≥14.3 ^g (9.85, 20.7)
	120 mg pibrentasvir (n=12)	300 mg QD with ritonavir 100 mg QD (n=12)	≥1.29 ^g (1.15, 1.45)	≥1.64 ^g (1.48, 1.82)	≥2.29 ^g (1.95, 2.68)
grazoprevir/elbasvir	grazoprevir 200 mg QD d 1–35 (n=12)	300 mg QD with ritonavir 100 mg QD d 1–35 (n=12)	6.24 (4.42, 8.81)	10.58 (7.78, 14.39)	11.64 (7.96, 17.02)
	elbasvir 50 mg QD d 1–35 (n=10)	300 mg QD with ritonavir 100 mg QD d 1–35 (n=10)	4.15 (3.46, 4.97)	4.76 (4.07, 5.56)	6.45 (5.51, 7.54)
methadone	Stable maintenance dose, d 1–15 (n=16)	400 mg QD, d 2–15 (n=16)	(R)-methadone ^h 0.91 (0.84, 1.0) total: 0.85 (0.78, 0.93)	(R)-methadone ^h 1.03 (0.95, 1.10) total: 0.94 (0.87, 1.02)	(R)-methadone ^h 1.11 (1.02, 1.20) total: 1.02 (0.93, 1.12)
nevirapine ^{ij}	200 mg BID, d 1–23 (n=23)	300 mg QD with ritonavir 100 mg QD, d 4–13, then 400 mg QD with ritonavir 100 mg QD, d 14–23 (n=23)	1.17 (1.09, 1.25) 1.21 (1.11, 1.32)	1.25 (1.17, 1.34) 1.26 (1.17, 1.36)	1.32 (1.22, 1.43) 1.35 (1.25, 1.47)

Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			C _{max}	AUC	C _{min}
omeprazole ^k	40 mg single dose, d 7 and d 20 (n=16)	400 mg QD, d 1–12 (n=16)	1.24 (1.04, 1.47)	1.45 (1.20, 1.76)	NA
rifabutin	300 mg QD, d 1–10 then 150 mg QD, d 11–20 (n=3)	600 mg QD, ^l d 11–20 (n=3)	1.18 (0.94, 1.48) 25-O-desacetyl-rifabutin: 8.20 (5.90, 11.40)	2.10 (1.57, 2.79) 25-O-desacetyl-rifabutin: 22.01 (15.97, 30.34)	3.43 (1.98, 5.96) 25-O-desacetyl-rifabutin: 75.6 (30.1, 190.0)
	150 mg twice weekly, d 1–15 (n=7)	300 mg QD with ritonavir 100 mg QD, d 1–17 (n=7)	2.49 ^m (2.03, 3.06) 25-O-desacetyl-rifabutin: 7.77 (6.13, 9.83)	1.48 ^m (1.19, 1.84) 25-O-desacetyl-rifabutin: 10.90 (8.14, 14.61)	1.40 ^m (1.05, 1.87) 25-O-desacetyl-rifabutin: 11.45 (8.15, 16.10)
pitavastatin	4 mg QD for 5 days	300 mg QD for 5 days	1.60 (1.39, 1.85)	1.31 (1.23, 1.39)	NA
rosiglitazone ⁿ	4 mg single dose, d 1, 7, 17 (n=14)	400 mg QD, d 2–7, then 300 mg QD with ritonavir 100 mg QD, d 8–17 (n=14)	1.08 (1.03, 1.13) 0.97 (0.91, 1.04)	1.35 (1.26, 1.44) 0.83 (0.77, 0.89)	NA NA
rosuvastatin	10 mg single dose	300 mg QD with ritonavir 100 mg QD for 7 days	↑ 7-fold ^o	↑ 3-fold ^o	NA
saquinavir ^p (soft gelatin capsules)	1200 mg QD, d 1–13 (n=7)	400 mg QD, d 7–13 (n=7)	4.39 (3.24, 5.95)	5.49 (4.04, 7.47)	6.86 (5.29, 8.91)
sofosbuvir/ velpatasvir/ voxilaprevir	400 mg sofosbuvir single dose (n=15)	300 mg with 100 mg ritonavir single dose (n=15)	1.29 (1.09, 1.52) sofosbuvir metabolite GS-331007 1.05 (0.99, 1.12)	1.40 (1.25, 1.57) sofosbuvir metabolite GS-331007 1.25 (1.16, 1.36)	NA
	100 mg velpatasvir single dose (n=15)	300 mg with 100 mg ritonavir single dose (n=15)	1.29 (1.07, 1.56)	1.93 (1.58, 2.36)	NA
	100 mg voxilaprevir single dose (n=15)	300 mg with 100 mg ritonavir single dose (n=15)	4.42 (3.65, 5.35)	4.31 (3.76, 4.93)	NA
tenofovir DF ^q	300 mg QD, d 9–16 (n=33) and d 24–30 (n=33)	400 mg QD, d 2–16 (n=33)	1.14 (1.08, 1.20)	1.24 (1.21, 1.28)	1.22 (1.15, 1.30)

Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ^a

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			C _{max}	AUC	C _{min}
	300 mg QD, d 1–7 (pm) (n=14) d 25–34 (pm) (n=12)	300 mg QD with ritonavir 100 mg QD, d 25–34 (am) (n=12) ^f	1.34 (1.20, 1.51)	1.37 (1.30, 1.45)	1.29 (1.21, 1.36)
voriconazole (Participants with at least one functional CYP2C19 allele)	200 mg BID, d 2–3, 22–30; 400 mg BID, d 1, 21 (n=20)	300 mg with ritonavir 100 mg QD, d 11–30 (n=20)	0.90 (0.78, 1.04)	0.67 (0.58, 0.78)	0.61 (0.51, 0.72)
voriconazole (Participants without a functional CYP2C19 allele)	50 mg BID, d 2–3, 22–30; 100 mg BID, d 1, 21 (n=8)	300 mg with ritonavir 100 mg QD, d 11–30 (n=8)	4.38 (3.55, 5.39)	5.61 (4.51, 6.99)	7.65 (5.71, 10.2)
lamivudine and zidovudine	150 mg lamivudine and 300 mg zidovudine BID, d 1–12 (n=19)	400 mg QD, d 7–12 (n=19)	lamivudine: 1.04 (0.92, 1.16) zidovudine: 1.05 (0.88, 1.24) zidovudine glucuronide: 0.95 (0.88, 1.02)	lamivudine: 1.03 (0.98, 1.08) zidovudine: 1.05 (0.96, 1.14) zidovudine glucuronide: 1.00 (0.97, 1.03)	lamivudine: 1.12 (1.04, 1.21) zidovudine: 0.69 (0.57, 0.84) zidovudine glucuronide: 0.82 (0.62, 1.08)

^a Data provided are under fed conditions unless otherwise noted.

^b 400 mg ddI EC and REYATAZ were administered together with food on Days 8 and 19.

^c Upon further dose normalization of ethinyl estradiol 25 mcg with atazanavir relative to ethinyl estradiol 35 mcg without atazanavir, the ratio of geometric means (90% confidence intervals) for C_{max}, AUC, and C_{min} were 0.82 (0.73, 0.92), 1.06 (0.95, 1.17), and 1.35 (1.11, 1.63), respectively.

^d Upon further dose normalization of ethinyl estradiol 35 mcg with atazanavir with ritonavir relative to ethinyl estradiol 25 mcg without atazanavir with ritonavir, the ratio of geometric means (90% confidence intervals) for C_{max}, AUC, and C_{min} were 1.17 (1.03, 1.34), 1.13 (1.05, 1.22), and 0.88 (0.77, 1.00), respectively.

^e All participants were on a 28-day lead-in period; one full cycle of Ortho Tri-Cyclen[®]. Ortho Tri-Cyclen[®] contains 35 mcg of ethinyl estradiol. Ortho Tri-Cyclen[®] LO contains 25 mcg of ethinyl estradiol. Results were dose normalized to an ethinyl estradiol dose of 35 mcg.

^f 17-deacetyl norgestimate is the active component of norgestimate.

^g Effect of atazanavir with ritonavir on the first dose of glecaprevir and pibrentasvir is reported.

^h (R)-methadone is the active isomer of methadone.

ⁱ Study was conducted in participants with HIV-1.

- ^j Participants were treated with nevirapine prior to study entry.
 - ^k Omeprazole was used as a metabolic probe for CYP2C19. Omeprazole was given 2 hours after REYATAZ on Day 7; and was given alone 2 hours after a light meal on Day 20.
 - ^l Not the recommended therapeutic dose of atazanavir.
 - ^m When compared to rifabutin 150 mg QD alone d1–10 (n=14). Total of rifabutin and 25-O-desacetyl-rifabutin: AUC 2.19 (1.78, 2.69).
 - ⁿ Rosiglitazone used as a probe substrate for CYP2C8.
 - ^o Mean ratio (with/without coadministered drug). ↑ indicates an increase in rosuvastatin exposure.
 - ^p The combination of atazanavir and saquinavir 1200 mg QD produced daily saquinavir exposures similar to the values produced by the standard therapeutic dosing of saquinavir at 1200 mg TID. However, the C_{max} is about 79% higher than that for the standard dosing of saquinavir (soft gelatin capsules) alone at 1200 mg TID.
 - ^q Note that similar results were observed in a study where administration of tenofovir DF and REYATAZ was separated by 12 hours.
 - ^r Administration of tenofovir DF and REYATAZ was temporally separated by 12 hours.
- NA = not available.

12.4 Microbiology

Mechanism of Action

Atazanavir (ATV) is an azapeptide HIV-1 protease inhibitor (PI). The compound selectively inhibits the virus-specific processing of viral Gag and Gag-Pol polyproteins in HIV-1-infected cells, thus preventing formation of mature virions.

Antiviral Activity in Cell Culture

Atazanavir exhibits anti-HIV-1 activity with a mean 50% effective concentration (EC_{50}) in the absence of human serum of 2 to 5 nM against a variety of laboratory and clinical HIV-1 isolates grown in peripheral blood mononuclear cells, macrophages, CEM-SS cells, and MT-2 cells. Atazanavir has activity against HIV-1 Group M subtype viruses A, B, C, D, AE, AG, F, G, and J isolates in cell culture. Atazanavir has variable activity against HIV-2 isolates (1.9–32 nM), with EC_{50} values above the EC_{50} values of failure isolates. Two-drug combination antiviral activity studies with atazanavir showed no antagonism in cell culture with PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir), NNRTIs (delavirdine, efavirenz, and nevirapine), NRTIs (abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir DF, and zidovudine), the HIV-1 fusion inhibitor enfuvirtide, and two compounds used in the treatment of viral hepatitis, adefovir and ribavirin, without enhanced cytotoxicity.

Resistance

In Cell Culture: HIV-1 isolates with a decreased susceptibility to atazanavir have been selected in cell culture and obtained from patients treated with atazanavir or atazanavir with ritonavir. HIV-1 isolates with 93- to 183-fold reduced susceptibility to atazanavir from three different viral strains were selected in cell culture by 5 months. The substitutions in these HIV-1 viruses that contributed to atazanavir resistance include I50L, N88S, I84V, A71V, and M46I. Changes were also observed

at the protease cleavage sites following drug selection. Recombinant viruses containing the I50L substitution without other major PI substitutions were growth impaired and displayed increased susceptibility in cell culture to other PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir). The I50L and I50V substitutions yielded selective resistance to atazanavir and amprenavir, respectively, and did not appear to be cross-resistant.

Clinical Studies of Treatment-Naive Participants: Comparison of Ritonavir-Boosted REYATAZ vs Unboosted REYATAZ: Study AI424-089 compared REYATAZ 300 mg once daily with ritonavir 100 mg vs REYATAZ 400 mg once daily when administered with lamivudine and extended-release stavudine in treatment-naive participants with HIV-1. A summary of the number of virologic failures and virologic failure isolates with atazanavir resistance in each arm is shown in Table 23.

Table 23: Summary of Virologic Failures^a at Week 96 in Study AI424-089: Comparison of Ritonavir Boosted REYATAZ vs Unboosted REYATAZ: Randomized Participants

	REYATAZ 300 mg with ritonavir 100 mg (n=95)	REYATAZ 400 mg (n=105)
Virologic Failure (≥50 copies/mL) at Week 96	15 (16%)	34 (32%)
Virologic Failure with Genotypes and Phenotypes Data	5	17
Virologic Failure Isolates with atazanavir - resistance at Week 96	0/5 (0%) ^b	4/17 (24%) ^b
Virologic Failure Isolates with I50L Emergence at Week 96 ^c	0/5 (0%) ^b	2/17 (12%) ^b
Virologic Failure Isolates with Lamivudine Resistance at Week 96	2/5 (40%) ^b	11/17 (65%) ^b

^a Virologic failure includes participants who were never suppressed through Week 96 and on study at Week 96, had virologic rebound or discontinued due to insufficient viral load response.

^b Percentage of Virologic Failure Isolates with genotypic and phenotypic data.

^c Mixture of I50I/L emerged in 2 other atazanavir 400 mg-treated participants. Neither isolate was phenotypically resistant to atazanavir.

Clinical Studies of Treatment-Naive Participants Receiving REYATAZ 300 mg with Ritonavir 100 mg: In Phase 3 Study AI424-138, an as-treated genotypic and phenotypic analysis was conducted on samples from participants who experienced virologic failure (HIV-1 RNA ≥400 copies/mL) or discontinued before achieving suppression on atazanavir with ritonavir (n=39; 9%) and lopinavir/ritonavir (n=39; 9%) through 96 weeks of treatment. In the atazanavir with ritonavir arm, one of the virologic failure isolates had a 56-fold decrease in atazanavir susceptibility emerge on therapy with the development of PI resistance-associated substitutions

L10F, V32I, K43T, M46I, A71I, G73S, I85I/V, and L90M. The NRTI resistance-associated substitution M184V also emerged on treatment in this isolate conferring emtricitabine resistance. Two atazanavir with ritonavir-virologic failure isolates had baseline phenotypic atazanavir resistance and IAS-defined major PI resistance-associated substitutions at baseline. The I50L substitution emerged on study in one of these failure isolates and was associated with a 17-fold decrease in atazanavir susceptibility from baseline and the other failure isolate with baseline atazanavir resistance and PI substitutions (M46M/I and I84I/V) had additional IAS-defined major PI substitutions (V32I, M46I, and I84V) emerge on atazanavir treatment associated with a 3-fold decrease in atazanavir susceptibility from baseline. Five of the treatment failure isolates in the atazanavir with ritonavir arm developed phenotypic emtricitabine resistance with the emergence of either the M184I (n=1) or the M184V (n=4) substitution on therapy and none developed phenotypic tenofovir disoproxil resistance. In the lopinavir/ritonavir arm, one of the virologic failure participant isolates had a 69-fold decrease in lopinavir susceptibility emerge on therapy with the development of PI substitutions L10V, V11I, I54V, G73S, and V82A in addition to baseline PI substitutions L10L/I, V32I, I54I/V, A71I, G73G/S, V82V/A, L89V, and L90M. Six lopinavir/ritonavir virologic failure isolates developed the M184V substitution and phenotypic emtricitabine resistance and two developed phenotypic tenofovir disoproxil resistance.

Clinical Studies of Treatment-Naive Participants Receiving REYATAZ 400 mg without Ritonavir: atazanavir-resistant clinical isolates from treatment-naive participants who experienced virologic failure on REYATAZ 400 mg treatment without ritonavir often developed an I50L substitution (after an average of 50 weeks of atazanavir therapy), often in combination with an A71V substitution, but also developed one or more other PI substitutions (eg, V32I, L33F, G73S, V82A, I85V, or N88S) with or without the I50L substitution. In treatment-naive participants, viral isolates that developed the I50L substitution, without other major PI substitutions, showed phenotypic resistance to atazanavir but retained in cell culture susceptibility to other PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir); however, there are no clinical data available to demonstrate the effect of the I50L substitution on the efficacy of subsequently administered PIs.

Clinical Studies of Treatment-Experienced Participants: In studies of treatment-experienced participants treated with atazanavir or atazanavir with ritonavir, most atazanavir-resistant isolates from participants who experienced virologic failure developed substitutions that were associated with resistance to multiple PIs and displayed decreased susceptibility to multiple PIs. The most common protease substitutions to develop in the viral isolates of participants who failed treatment with atazanavir 300 mg once daily and ritonavir 100 mg once daily (together with tenofovir DF and an NRTI) included V32I, L33F/V/I, E35D/G, M46I/L, I50L, F53L/V, I54V, A71V/T/I, G73S/T/C, V82A/T/L, I85V, and L89V/Q/M/T. Other substitutions that developed on atazanavir with ritonavir treatment including E34K/A/Q, G48V, I84V, N88S/D/T, and L90M occurred in less than 10% of participant isolates. Generally, if multiple PI resistance substitutions were present in the HIV-1 virus of the participant at baseline, atazanavir resistance developed through substitutions associated with resistance to other PIs and could include the development of the I50L substitution. The I50L substitution has been detected in treatment-experienced participants experiencing

virologic failure after long-term treatment. Protease cleavage site changes also emerged on atazanavir treatment, but their presence did not correlate with the level of atazanavir resistance.

Clinical Studies of Pediatric Participants in AI424-397 (PRINCE I) and AI424-451 (PRINCE II): Treatment-emergent atazanavir with ritonavir resistance-associated amino acid substitution M36I in the protease was detected in the virus of one participant among treatment failures in AI424-397. In addition, three known resistance-associated substitutions for other PIs arose in the viruses from one participant each (L19I/R, H69K/R, and I72I/V). Reduced susceptibility to atazanavir, ritonavir, or atazanavir with ritonavir was not seen with these viruses. In AI424-451, atazanavir with ritonavir resistance-associated substitutions G16E, V82A/I/T, I84V, and/or L90M arose in the viruses of two participants. The virus population harboring the M46M/V, V82V/I, I84I/V, and L90L/M substitutions acquired phenotypic resistance to ritonavir (ritonavir phenotypic fold-change of 3.5, with a ritonavir cutoff of 2.5-fold change). However, these substitutions did not result in phenotypic resistance to atazanavir (atazanavir phenotypic fold-change of <1.8, with an atazanavir cutoff of 2.2-fold change). Secondary PI resistance-associated amino acid substitutions also arose in the viruses of one participant each, including V11V/I, D30D/G, E35E/D, K45K/R, L63P/S, and I72I/T. Q61D and Q61E/G emerged in the viruses of two participants who failed treatment with atazanavir with ritonavir. Viruses from nine participants in the two studies developed NRTI resistance-associated substitutions: K65K/R (n=1), M184V (n=7), and T215I (n=1).

Cross-Resistance

Cross-resistance among PIs has been observed. Baseline phenotypic and genotypic analyses of clinical isolates from atazanavir clinical trials of PI-experienced participants showed that isolates cross-resistant to multiple PIs were cross-resistant to atazanavir. Greater than 90% of the isolates with substitutions that included I84V or G48V were resistant to atazanavir. Greater than 60% of isolates containing L90M, G73S/T/C, A71V/T, I54V, M46I/L, or a change at V82 were resistant to atazanavir, and 38% of isolates containing a D30N substitution in addition to other changes were resistant to atazanavir. Isolates resistant to atazanavir were also cross-resistant to other PIs with >90% of the isolates resistant to indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir, and 80% resistant to amprenavir. In treatment-experienced participants, PI-resistant viral isolates that developed the I50L substitution in addition to other PI resistance-associated substitution were also cross-resistant to other PIs.

Baseline Genotype/Phenotype and Virologic Outcome Analyses

Genotypic and/or phenotypic analysis of baseline virus may aid in determining atazanavir susceptibility before initiation of atazanavir with ritonavir therapy. An association between virologic response at 48 weeks and the number and type of primary PI resistance-associated substitutions detected in baseline HIV-1 isolates from antiretroviral-experienced participants receiving atazanavir with ritonavir once daily or lopinavir / ritonavir (fixed-dose product) twice daily in Study AI424-045 is shown in Table 24.

Overall, both the number and type of baseline PI substitutions affected response rates in treatment-experienced participants. In the atazanavir with ritonavir group, participants had lower response

rates when 3 or more baseline PI substitutions, including a substitution at position 36, 71, 77, 82, or 90, were present compared to participants with 1–2 PI substitutions, including one of these substitutions.

Table 24: HIV-1 RNA Response by Number and Type of Baseline PI Substitution, Antiretroviral-Experienced Participants in Study AI424-045, As-Treated Analysis

Number and Type of Baseline PI Substitutions ^a	Virologic Response = HIV RNA <400 copies/mL ^b	
	atazanavir with ritonavir (n=110)	lopinavir/ritonavir ^c (n=113)
3 or more primary PI substitutions including^d:		
D30N	75% (6/8)	50% (3/6)
M36I/V	19% (3/16)	33% (6/18)
M46I/L/T	24% (4/17)	23% (5/22)
I54V/L/T/M/A	31% (5/16)	31% (5/16)
A71V/T/I/G	34% (10/29)	39% (12/31)
G73S/A/C/T	14% (1/7)	38% (3/8)
V77I	47% (7/15)	44% (7/16)
V82A/F/T/S/I	29% (6/21)	27% (7/26)
I84V/A	11% (1/9)	33% (2/6)
N88D	63% (5/8)	67% (4/6)
L90M	10% (2/21)	44% (11/25)
Number of baseline primary PI substitutions^a		
All patients, as-treated	58% (64/110)	59% (67/113)
0–2 PI substitutions	75% (50/67)	75% (50/67)
3–4 PI substitutions	41% (14/34)	43% (12/28)
5 or more PI substitutions	0% (0/9)	28% (5/18)

^a Primary substitutions include any change at D30, V32, M36, M46, I47, G48, I50, I54, A71, G73, V77, V82, I84, N88, and L90.

^b Results should be interpreted with caution because the subgroups were small.

^c Administered as a fixed-dose product.

^d There were insufficient data (n<3) for PI substitutions V32I, I47V, G48V, I50V, and F53L.

The response rates of antiretroviral-experienced participants in Study AI424-045 were analyzed by baseline phenotype (shift in susceptibility in cell culture relative to reference, Table 25). The analyses are based on a select population with 62% of participants receiving an NNRTI-based

regimen before study entry compared to 35% receiving a PI-based regimen. Additional data are needed to determine clinically relevant break points for REYATAZ.

Table 25: Baseline Phenotype by Outcome, Antiretroviral-Experienced Participants in Study AI424-045, As-Treated Analysis

Baseline Phenotype ^a	Virologic Response = HIV-1 RNA <400 copies/mL ^b	
	atazanavir with ritonavir (n=111)	lopinavir/ritonavir ^c (n=111)
0–2	71% (55/78)	70% (56/80)
>2–5	53% (8/15)	44% (4/9)
>5–10	13% (1/8)	33% (3/9)
>10	10% (1/10)	23% (3/13)

^a Fold change susceptibility in cell culture relative to the wild-type reference.

^b Results should be interpreted with caution because the subgroups were small.

^c Administered as a fixed-dose product.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term carcinogenicity studies in mice and rats were carried out with atazanavir for two years. In the mouse study, drug-related increases in hepatocellular adenomas were found in females at 360 mg/kg/day. The systemic drug exposure (AUC) at the NOAEL (no observable adverse effect level) in females, (120 mg/kg/day) was 2.8 times and in males (80 mg/kg/day) was 2.9 times higher than those in humans at the clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir, non-pregnant patients). In the rat study, no drug-related increases in tumor incidence were observed at doses up to 1200 mg/kg/day, for which AUCs were 1.1 (males) or 3.9 (females) times those measured in humans at the clinical dose.

Mutagenesis

Atazanavir tested positive in an *in vitro* clastogenicity test using primary human lymphocytes, in the absence and presence of metabolic activation. Atazanavir tested negative in the *in vitro* Ames reverse-mutation assay, *in vivo* micronucleus and DNA repair tests in rats, and *in vivo* DNA damage test in rat duodenum (comet assay).

Impairment of Fertility

At the systemic drug exposure levels (AUC) 0.9 (in male rats) or 2.3 (in female rats) times that of the human clinical dose, (300 mg/day atazanavir boosted with 100 mg/day ritonavir) significant effects on mating, fertility, or early embryonic development were not observed.

14 CLINICAL STUDIES

14.1 Adult Participants without Prior Antiretroviral Therapy

Study AI424-138: a 96-week study comparing the antiviral efficacy and safety of either REYATAZ or lopinavir/ritonavir, each in combination with fixed-dose tenofovir DF/emtricitabine in treatment-naïve participants with HIV-1 infection. Study AI424-138 (NCT00272779) was a 96-week, open-label, randomized, multicenter study, comparing REYATAZ (300 mg once daily) with ritonavir (100 mg once daily) to lopinavir/ritonavir (400/100 mg twice daily as fixed-dose product), each in combination with the fixed-dose product, tenofovir DF/emtricitabine (300/200 mg once daily), in 878 antiretroviral treatment-naïve participants. Participants had a mean age of 36 years (range: 19-72), 49% were Caucasian, 18% Black, 9% Asian, 23% Hispanic/Mestizo/mixed race, and 68% were male. The median baseline plasma CD4⁺ cell count was 204 cells/mm³ (range: 2 to 810 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.94 log₁₀ copies/mL (range: 2.60 to 5.88 log₁₀ copies/mL). Treatment response and outcomes through Week 96 are presented in Table 26.

Table 26: Outcomes of Treatment Through Week 96 in Treatment-Naïve Adults (Study AI424-138)

Outcome	REYATAZ 300 mg with ritonavir 100 mg (once daily) and tenofovir DF/emtricitabine (once daily) ^a (n=441)	lopinavir/ritonavir ^b 400 mg/100 mg (twice daily) with tenofovir DF/emtricitabine (once daily) ^a (n=437)
	96 Weeks	96 Weeks
Responder ^{c,d,e}	75%	68%
Virologic failure ^f	17%	19%
Rebound	8%	10%
Never suppressed through Week 96	9%	9%
Death	1%	1%
Discontinued due to adverse event	3%	5%
Discontinued for other reasons ^g	4%	7%

^a As a fixed-dose product: 300 mg tenofovir DF/200 mg emtricitabine once daily.

^b As a fixed-dose product: 400 mg lopinavir/100 mg ritonavir (twice daily).

^c Participants achieved HIV-1 RNA <50 copies/mL at Week 96. Roche Amplicor®, v1.5 ultra-sensitive assay.

^d Pre-specified ITT analysis at Week 48 using as-randomized cohort: atazanavir with ritonavir 78% and lopinavir/ritonavir 76% (difference estimate: 1.7% [95% confidence interval: -3.8%, 7.1%]).

^e Pre-specified ITT analysis at Week 96 using as-randomized cohort: atazanavir with ritonavir 74% and lopinavir/ritonavir 68% (difference estimate: 6.1% [95% confidence interval: 0.3%, 12.0%]).

- ^f Includes viral rebound and failure to achieve confirmed HIV-1 RNA <50 copies/mL through Week 96.
- ^g Includes lost to follow-up, participant's withdrawal, noncompliance, protocol violation, and other reasons.

Through 96 weeks of therapy, the proportion of responders among participants with high viral loads (ie, baseline HIV-1 RNA $\geq 100,000$ copies/mL) was comparable for the REYATAZ with ritonavir (165 of 223 participants, 74%) and lopinavir/ritonavir (148 of 222 participants, 67%) arms. At 96 weeks, the median increase from baseline in CD4+ cell count was 261 cells/mm³ for the REYATAZ with ritonavir arm and 273 cells/mm³ for the lopinavir/ritonavir arm.

Study AI424-034: REYATAZ once daily compared to efavirenz once daily, each in combination with fixed-dose lamivudine/zidovudine twice daily. Study AI424-034 (NCT00013897) was a randomized, double-blind, multicenter trial comparing REYATAZ (400 mg once daily) to efavirenz (600 mg once daily), each in combination with the fixed-dose product of lamivudine/zidovudine (150 mg/300 mg) given twice daily, in 810 antiretroviral treatment-naïve participants. Participants had a mean age of 34 years (range: 18 to 73), 36% were Hispanic, 33% were Caucasian, and 65% were male. The mean baseline CD4+ cell count was 321 cells/mm³ (range: 64 to 1424 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.8 log₁₀ copies/mL (range: 2.2 to 5.9 log₁₀ copies/mL). Treatment response and outcomes through Week 48 are presented in Table 27.

Table 27: Outcomes of Randomized Treatment Through Week 48 in Treatment-Naïve Adults (Study AI424-034)

Outcome	REYATAZ 400 mg once daily and lamivudine/ zidovudine ^d (n=405)	efavirenz 600 mg once daily and lamivudine/ zidovudine ^d (n=405)
Responder ^a	67% (32%)	62% (37%)
Virologic failure ^b	20%	21%
Rebound	17%	16%
Never suppressed through Week 48	3%	5%
Death	—	<1%
Discontinued due to adverse event	5%	7%
Discontinued for other reasons ^c	8%	10%

^a Participants achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48. Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.0 or 1.5 as geographically appropriate.

^b Includes viral rebound and failure to achieve confirmed HIV-1 RNA <400 copies/mL through Week 48.

^c Includes lost to follow-up, participant's withdrawal, noncompliance, protocol violation, and other reasons.

^d As a fixed-dose product: 150 mg lamivudine/300 mg zidovudine twice daily.

Through 48 weeks of therapy, the proportion of responders among participants with high viral loads (ie, baseline HIV-1 RNA $\geq 100,000$ copies/mL) was comparable for the REYATAZ and efavirenz arms. The mean increase from baseline in CD4+ cell count was 176 cells/mm³ for the REYATAZ arm and 160 cells/mm³ for the efavirenz arm.

Study AI424-008: REYATAZ 400 mg once daily compared to REYATAZ 600 mg once daily, and compared to nelfinavir 1250 mg twice daily, each in combination with stavudine and lamivudine twice daily. Study AI424-008 (NCT identifier not available) was a 48-week, randomized, multicenter trial, blinded to dose of REYATAZ, comparing REYATAZ at two dose levels (400 mg and 600 mg once daily) to nelfinavir (1250 mg twice daily), each in combination with stavudine (40 mg) and lamivudine (150 mg) given twice daily, in 467 antiretroviral treatment-naïve participants. Participants had a mean age of 35 years (range: 18 to 69), 55% were Caucasian, and 63% were male. The mean baseline CD4+ cell count was 295 cells/mm³ (range: 4 to 1003 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.7 log₁₀ copies/mL (range: 1.8 to 5.9 log₁₀ copies/mL). Treatment response and outcomes through Week 48 are presented in Table 28.

Table 28: Outcomes of Randomized Treatment Through Week 48 in Treatment-Naïve Adults (Study AI424-008)

Outcome	REYATAZ 400 mg once daily with lamivudine and stavudine (n=181)	nelfinavir 1250 mg twice daily with lamivudine and stavudine (n=91)
Responder ^a	67% (33%)	59% (38%)
Virologic failure ^b	24%	27%
Rebound	14%	14%
Never suppressed through Week 48	10%	13%
Death	<1%	—
Discontinued due to adverse event	1%	3%
Discontinued for other reasons ^c	7%	10%

^a Participants achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48. Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.0 or 1.5 as geographically appropriate.

^b Includes viral rebound and failure to achieve confirmed HIV-1 RNA <400 copies/mL through Week 48.

^c Includes lost to follow-up, participant's withdrawal, noncompliance, protocol violation, and other reasons.

Through 48 weeks of therapy, the mean increase from baseline in CD4+ cell count was 234 cells/mm³ for the REYATAZ 400-mg arm and 211 cells/mm³ for the nelfinavir arm.

14.2 Adult Participants with Prior Antiretroviral Therapy

Study AI424-045: REYATAZ once daily with ritonavir once daily compared to REYATAZ once daily and saquinavir (soft gelatin capsules) once daily, and compared to lopinavir/ritonavir twice daily, each in combination with tenofovir DF and one NRTI. Study AI424-045 (NCT00035932): was a randomized, multicenter trial comparing REYATAZ (300 mg once daily) with ritonavir (100 mg once daily) to REYATAZ (400 mg once daily) with saquinavir soft gelatin capsules (1200 mg once daily), and to lopinavir/ritonavir (400/100 mg twice daily as fixed-dose product), each in combination with tenofovir DF and one NRTI, in 347 (of 358 randomized) participants who experienced virologic failure on highly active antiretroviral therapy regimens containing PIs, NNRTIs, and NRTIs. The mean time of prior exposure to antiretrovirals was 139 weeks for PIs, 85 weeks for NNRTIs, and 283 weeks for NRTIs. The mean age was 41 years (range: 24 to 74); 60% were Caucasian, and 78% were male. The mean baseline CD4⁺ cell count was 338 cells/mm³ (range: 14 to 1543 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.4 log₁₀ copies/mL (range: 2.6 to 5.88 log₁₀ copies/mL).

Treatment outcomes through Week 48 for the REYATAZ with ritonavir and lopinavir/ritonavir treatment arms are presented in Table 29. REYATAZ with ritonavir and lopinavir/ritonavir were similar for the primary efficacy outcome measure of time-averaged difference in change from baseline in HIV-1 RNA level. Study AI424-045 was not large enough to reach a definitive conclusion that REYATAZ with ritonavir and lopinavir/ritonavir are equivalent on the secondary efficacy outcome measure of proportions below the HIV-1 RNA lower limit of quantification [*see Microbiology, Tables 24 and 25 (12.4)*].

Table 29: Outcomes of Treatment Through Week 48 in Study AI424-045 (Participants with Prior Antiretroviral Experience)

Outcome	REYATAZ 300 mg with ritonavir 100 mg once daily and tenofovir DF and 1 NRTI (n=119)	lopinavir/ritonavir (400/100 mg) twice daily and tenofovir DF and 1 NRTI (n=118)	Difference ^a (REYATAZ- lopinavir/ritonavir) ^b (CI)
HIV-1 RNA Change from Baseline (log ₁₀ copies/mL) ^c	-1.58	-1.70	+0.12 ^c (-0.17, 0.41)
CD4+ Change from Baseline (cells/mm ³) ^c	116	123	-7 (-67, 52)
Percent of Participants Responding ^c			
HIV-1 RNA <400 copies/mL ^c	55%	57%	-2.2% (-14.8%, 10.5%)
HIV-1 RNA <50 copies/mL ^c	38%	45%	-7.1% (-19.6%, 5.4%)

^a Time-averaged difference through Week 48 for HIV-1 RNA; Week 48 difference in HIV-1 RNA percentages and CD4+ mean changes, REYATAZ with ritonavir vs lopinavir/ritonavir; CI = 97.5% confidence interval for change in HIV-1 RNA; 95% confidence interval otherwise.

^b Administered as a fixed-dose product.

^c Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.5.

^d Protocol-defined primary efficacy outcome measure.

^e Based on participants with baseline and Week 48 CD4+ cell count measurements (REYATAZ with ritonavir, n=85; lopinavir/ritonavir, n=93).

^f Participants achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48.

No participants in the REYATAZ with ritonavir treatment arm and three participants in the lopinavir/ritonavir treatment arm experienced a new-onset CDC Category C event during the study.

In Study AI424-045, the mean change from baseline in plasma HIV-1 RNA for REYATAZ 400 mg with saquinavir (n=115) was -1.55 log₁₀ copies/mL, and the time-averaged difference in change in HIV-1 RNA levels versus lopinavir/ritonavir was 0.33. The corresponding mean increase in CD4+ cell count was 72 cells/mm³. Through 48 weeks of treatment, the proportion of participants in this treatment arm with plasma HIV-1 RNA <400 (<50) copies/mL was 38% (26%). In this study, coadministration of REYATAZ and saquinavir did not provide adequate efficacy [see *Drug Interactions (7)*].

Study AI424-045 also compared changes from baseline in lipid values. [See *Adverse Reactions* (6.1).]

Study AI424-043 (NCT00028301): Study AI424-043 was a randomized, open-label, multicenter trial comparing REYATAZ (400 mg once daily) to lopinavir/ritonavir (400/100 mg twice daily as fixed-dose product), each in combination with two NRTIs, in 300 participants who experienced virologic failure to only one prior PI-containing regimen. Through 48 weeks, the proportion of participants with plasma HIV-1 RNA <400 (<50) copies/mL was 49% (35%) for participants randomized to REYATAZ (n=144) and 69% (53%) for participants randomized to lopinavir/ritonavir (n=146). The mean change from baseline was $-1.59 \log_{10}$ copies/mL in the REYATAZ treatment arm and $-2.02 \log_{10}$ copies/mL in the lopinavir/ritonavir arm. Based on the results of this study, REYATAZ without ritonavir was inferior to lopinavir/ritonavir in PI-experienced participants with prior virologic failure and is not recommended for such patients.

14.3 Pediatric Participants

Pediatric Trials with REYATAZ Capsules

Study AI424-040; PACTG 1020A (NCT00006604): Assessment of the pharmacokinetics, safety, tolerability, and virologic response of REYATAZ capsules was based on data from this open-label, multicenter clinical trial which included participants from 6 years to 21 years of age. In this study, 105 participants (43 antiretroviral-naïve and 62 antiretroviral-experienced) received once daily REYATAZ capsule formulation, with or without ritonavir, in combination with two NRTIs.

One-hundred five (105) participants (6 to less than 18 years of age) treated with the REYATAZ capsule formulation, with or without ritonavir, were evaluated. Using an intent-to-treat (ITT) analysis, the overall proportions of antiretroviral-naïve and -experienced participants with HIV-1 RNA <400 copies/mL at Week 96 were 51% (22/43) and 34% (21/62), respectively. The overall proportions of antiretroviral-naïve and -experienced participants with HIV-1 RNA <50 copies/mL at Week 96 were 47% (20/43) and 24% (15/62), respectively. The median increase from baseline in absolute CD4 count at 96 weeks of therapy was 335 cells/mm³ in antiretroviral-naïve participants and 220 cells/mm³ in antiretroviral-experienced participants.

Pediatric Trials with REYATAZ Oral Powder

Assessment of the pharmacokinetics, safety, tolerability, and virologic response of REYATAZ oral powder was based on data from two open-label, multicenter clinical trials.

- AI424-397 (PRINCE I; *NCT01099579*): In pediatric participants from 3 months to less than 6 years of age
- AI424-451 (PRINCE II; *NCT01335698*): In pediatric participants from 3 months to less than 11 years of age

In these studies, 155 participants (59 antiretroviral-naïve and 96 antiretroviral-experienced) received once daily REYATAZ oral powder with ritonavir, in combination with two NRTIs.

For inclusion in both trials, treatment-naïve participants had to have genotypic sensitivity to REYATAZ and two NRTIs, and treatment-experienced participants had to have documented

genotypic and phenotypic sensitivity at screening to REYATAZ and at least 2 NRTIs. Participants exposed only to antiretrovirals *in utero* or intrapartum were considered treatment-naïve. Participants who received REYATAZ or REYATAZ with ritonavir at any time prior to study enrollment or who had a history of treatment failure on two or more protease inhibitors were excluded from the trials.

One hundred thirty-four (134) participants from both studies weighing 5 kg to less than 35 kg treated with REYATAZ oral powder with ritonavir were evaluated. Participants 5 kg to less than 10 kg received either 150 mg or 200 mg REYATAZ and 80 mg ritonavir oral solution; participants 10 kg to less than 15 kg received 200 mg REYATAZ and 80 mg ritonavir oral solution; participants 15 kg to less than 25 kg received 250 mg REYATAZ and 80 mg ritonavir oral solution; and participants 25 kg to less than 35 kg received 300 mg REYATAZ and 100 mg ritonavir.

Using a modified ITT analysis, the overall proportions of antiretroviral-naïve and antiretroviral-experienced participants with HIV-1 RNA <400 copies/mL at Week 48 were 79% (41/52) and 62% (51/82), respectively in participants who received REYATAZ oral powder with ritonavir. The overall proportions of antiretroviral-naïve and antiretroviral-experienced participants with HIV-1 RNA <50 copies/mL at Week 48 were 54% (28/52) and 50% (41/82), respectively, in participants who received REYATAZ oral powder with ritonavir. The median increase from baseline in absolute CD4 count (percent) at 48 weeks of therapy (last observation carried forward) was 215 cells/mm³ (6%) in antiretroviral-naïve participants and 133 cells/mm³ (4%) in antiretroviral-experienced participants who received REYATAZ oral powder with ritonavir.

16 HOW SUPPLIED/STORAGE AND HANDLING

REYATAZ Capsules

REYATAZ[®] (atazanavir) capsules are available in the following strengths and configurations of plastic bottles with child-resistant closures.

Product Strength*	Capsule Shell Color (cap/body)	Markings on Capsule (ink color)		Capsules per Bottle	NDC Number
		cap	body		
200 mg	blue/blue	BMS 200 mg (white)	3631 (white)	60	0003-3631-12
300 mg	red/blue	BMS 300 mg (white)	3622 (white)	30	0003-3622-12

* 200 mg atazanavir equivalent to 227.8 mg atazanavir sulfate.

300 mg atazanavir equivalent to 341.69 mg atazanavir sulfate.

Keep capsules in a tightly closed container.

Store REYATAZ capsules at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

REYATAZ Oral Powder

REYATAZ oral powder is an orange-vanilla flavored powder, packed in child-resistant packets. Each packet contains 50 mg of atazanavir equivalent to 56.9 mg of atazanavir sulfate in 1.5 g of powder. REYATAZ oral powder is supplied in cartons (NDC 0003-3638-10) of 30 packets each. [See *Dosage and Administration* (2.5)].

Store REYATAZ oral powder at a temperature of 68°F to 86°F (20°C to 30°C). Store REYATAZ oral powder in the original packet. Do not open until ready to use. After REYATAZ oral powder is mixed with food or liquid, it may be kept at a temperature 68°F to 86°F (20°C to 30°C) for up to 1 hour. Take REYATAZ oral powder within 1 hour after mixing with food or liquid.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (*Patient Information and Instructions for Use*).

REYATAZ is not a cure for HIV-1 infection. Advise patients to remain under the care of a healthcare provider while using REYATAZ.

Cardiac Conduction Abnormalities

Inform patients that atazanavir may produce changes in the electrocardiogram (eg, PR prolongation). Tell patients to consult their healthcare provider if they are experiencing symptoms such as dizziness or lightheadedness [see *Warnings and Precautions* (5.1)].

Severe Skin Reaction

Inform patients that there have been reports of severe skin reactions (eg, Stevens-Johnson syndrome, erythema multiforme, and toxic skin eruptions) with REYATAZ use. Advise patients that if signs or symptoms of severe skin reactions or hypersensitivity reactions develop, they must discontinue REYATAZ and seek medical evaluation immediately [see *Warnings and Precautions* (5.2) and *Adverse Reactions* (6.1)].

Hyperbilirubinemia

Inform patients that asymptomatic elevations in indirect bilirubin have occurred in patients receiving REYATAZ. This may be accompanied by yellowing of the skin or whites of the eyes and alternative antiretroviral therapy may be considered if the patient has cosmetic concerns [see *Warnings and Precautions* (5.8)].

Patients with Phenylketonuria

Advise caregivers of patients with phenylketonuria that REYATAZ oral powder contains phenylalanine [see *Warnings and Precautions* (5.3)].

Chronic Kidney Disease

Inform patients that treatment with REYATAZ may lead to the development of chronic kidney disease, and to maintain adequate hydration while taking REYATAZ [see *Warnings and Precautions* (5.5)].

Nephrolithiasis and Cholelithiasis

Inform patients that kidney stones and/or gallstones have been reported with REYATAZ use. Some patients with kidney stones and/or gallstones required hospitalization for additional management, and some had complications. Discontinuation of REYATAZ may be necessary as part of the medical management of these adverse events [see *Warnings and Precautions* (5.6)].

Drug Interactions

REYATAZ may lead to significant interaction with some drugs; therefore, advise patients to report the use of any other prescription, nonprescription medication, or herbal products, particularly St. John's wort, to their healthcare provider prior to use [see *Contraindications* (4), *Warnings and Precautions* (5.7)].

Immune Reconstitution Syndrome

Advise patients to inform their healthcare provider immediately of any symptoms of infection, as in some patients with advanced HIV-1, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV-1 treatment is started [see *Warnings and Precautions* (5.10)].

Fat Redistribution

Inform patients that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy including protease inhibitors and that the cause and long-term health effects of these conditions are not known at this time [see *Warnings and Precautions* (5.11)].

Dosing Instructions

Advise patients to take REYATAZ with food every day and take other concomitant antiretroviral therapy as prescribed. REYATAZ must always be used in combination with other antiretroviral drugs. Advise patients that they should not alter the dose or discontinue therapy without consulting with their healthcare provider. Tell patients if a dose of REYATAZ is missed, they should take the dose as soon as possible and then return to their normal schedule; however, if a dose is skipped the patient should not double the next dose.

Pregnancy

Inform pregnant patients that there is a pregnancy exposure registry that monitors pregnancy outcomes in pregnant patients exposed to REYATAZ during pregnancy. Healthcare providers are encouraged to register patients by calling the Antiretroviral Pregnancy Registry [see *Use in Specific Populations* (8.1)].

Lactation

Instruct patients with HIV-1 that the potential risks of breastfeeding include: (1) HIV-1 transmission to infants without HIV-1, (2) developing viral resistance in infants with HIV-1, and (3) adverse reactions in a breastfed infant similar to those seen in adults [see *Use in Specific Populations* (8.2)].

PATIENT INFORMATION	
REYATAZ® (RAY-ah-taz) (atazanavir) capsules	REYATAZ® (RAY-ah-taz) (atazanavir) oral powder
Important: Ask your healthcare provider or pharmacist about medicines that should not be taken with REYATAZ. For more information, see “Do not take REYATAZ if you” and “Before taking REYATAZ”.	
What is REYATAZ? REYATAZ is a prescription medicine that is used to treat human immunodeficiency virus-1 (HIV-1) infection, in combination with other HIV-1 medicines in adults and children 3 months of age and older and who weigh at least 11 pounds (5 kg). HIV-1 is the virus that causes AIDS (Acquired Immunodeficiency Syndrome). REYATAZ should not be used in children younger than 3 months of age.	
Do not take REYATAZ if you: - are allergic to atazanavir or any of the ingredients in REYATAZ. See the end of this leaflet for a complete list of ingredients in REYATAZ. - are taking any of the following medicines. Taking REYATAZ with these medicines may affect how REYATAZ works. REYATAZ may cause serious or life-threatening side effects, or death when used with these medicines: - alfuzosin - amiodarone (when REYATAZ is used with ritonavir) - apalutamide - carbamazepine - cisapride - elbasvir and grazoprevir - encorafenib - ergot medicines including: - dihydroergotamine - ergonovine - ergonovine ergotamine - methylergonovine - glecaprevir and pibrentasvir - indinavir - irinotecan - ivosidenib - lurasidone (when REYATAZ is used with ritonavir) - lomitapide - lovastatin - midazolam, when taken by mouth for sedation - nevirapine - phenobarbital - phenytoin - pimozide - quinidine (when REYATAZ is used with ritonavir) - rifampin - sildenafil, when used for the treatment of pulmonary arterial hypertension - simvastatin - St. John's wort - triazolam	
Before taking REYATAZ, tell your healthcare provider about all of your medical conditions, including if you: - have heart problems - have liver problems, including hepatitis B or C virus - have phenylketonuria (PKU). The artificial sweetener aspartame in REYATAZ oral powder contains phenylalanine, which can be harmful to people with PKU. - have kidney problems - are receiving dialysis treatment - have diabetes - have hemophilia - are pregnant or plan to become pregnant. REYATAZ must be taken with ritonavir during pregnancy. Hormonal forms of birth control, such as injections, vaginal rings or implants, contraceptive patch, and some birth control pills may not work during treatment with REYATAZ. Talk to your healthcare provider about forms of birth control that may be used during treatment with REYATAZ. Pregnancy Exposure Registry. There is a pregnancy exposure registry for people who take REYATAZ during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.	

- **After your baby is born**, tell your healthcare provider if your baby's skin or the white part of their eyes turns yellow.
- are breastfeeding or plan to breastfeed. REYATAZ can pass into your breast milk.
- Talk to your healthcare provider about the following risks of breastfeeding during treatment with REYATAZ:
 - The HIV-1 virus may pass to your baby if your baby does not have the HIV-1 virus.
 - The HIV-1 virus may become harder to treat if your baby has the HIV-1 virus.
 - Your baby may get side effects from REYATAZ.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Some medicines interact with REYATAZ. **Keep a list of your medicines to show your healthcare provider and pharmacist.**

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with REYATAZ.
- **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take REYATAZ with other medicines.

How should I take REYATAZ?

- **Take REYATAZ exactly as your healthcare provider tells you to.**
- Do not change your dose or stop taking REYATAZ unless your healthcare provider tells you to.
- Stay under the care of your healthcare provider during treatment with REYATAZ.
- REYATAZ must be used with other HIV-1 medicines.
- Take REYATAZ 1 time each day.
- REYATAZ comes as capsules and oral powder.
- Take REYATAZ capsules and oral powder with food.
- Swallow the capsules whole. Do not open the capsules.
- REYATAZ oral powder must be mixed with food or liquid. Your child's healthcare provider will prescribe the right dose of REYATAZ based on your child's weight. **See the detailed "Instructions for Use" that comes with REYATAZ oral powder for information about the correct way to mix and give a dose of REYATAZ oral powder to your child.**
- REYATAZ oral powder must be taken with ritonavir.
- If you miss a dose of REYATAZ, take it as soon as you remember. Then take the next dose at your regular time. Do not take 2 doses at the same time.
- If you take too much REYATAZ, call your healthcare provider or go to the nearest hospital emergency room right away.

When your supply of REYATAZ starts to run low, get more from your healthcare provider or pharmacy. It is important not to run out of REYATAZ. The amount of HIV-1 in your blood may increase if the medicine is stopped for even a short time. The virus may become resistant to REYATAZ and harder to treat.

What are the possible side effects of REYATAZ?

REYATAZ can cause serious side effects, including:

- **A change in the way your heart beats (heart rhythm change).** Tell your healthcare provider right away if you get dizzy or lightheaded. These could be symptoms of a heart problem.
- **Skin rash.** Skin rash is common with REYATAZ but can sometimes be severe. Severe rash may develop with other symptoms which could be serious. If you develop a severe rash or a rash with any of the following symptoms, stop taking REYATAZ and call your healthcare provider or go to the nearest hospital emergency room right away:

○ general feeling of discomfort or "flu-like" symptoms	○ blisters
○ fever	○ mouth sores
○ muscle or joint aches	○ swelling of your face
○ red or inflamed eyes, like "pink eye" (conjunctivitis)	○ painful, warm, or red lump under your skin
- **Liver problems.** If you have liver problems, including hepatitis B or C virus, your liver problems may get worse when you take REYATAZ. Your healthcare provider will do blood tests to check your liver before you start REYATAZ and during treatment. Tell your healthcare provider right away if you get any of the following symptoms:

○ dark "tea-colored" urine	○ nausea
○ your skin or the white part of your eyes turns yellow	○ itching
○ light colored stools	○ stomach-area pain

- **Chronic kidney disease.** REYATAZ may affect how well your kidneys work. Your healthcare provider will do blood and urine tests to check your kidneys before you start REYATAZ and during treatment. Drink plenty of fluids during treatment with REYATAZ.
- **Kidney stones** have happened in some people who take REYATAZ, and sometimes may lead to hospitalization. Tell your healthcare provider right away if you get symptoms of kidney stones which may include pain in your low back or low stomach area, blood in your urine, or pain when you urinate.
- **Gallbladder stones** have happened in some people who take REYATAZ, and sometimes may lead to hospitalization. Tell your healthcare provider right away if you get symptoms of a gallbladder problem which may include:
 - pain in the right or middle upper stomach area
 - nausea and vomiting
 - fever
 - your skin or the white part of your eyes turns yellow
- **Yellowing of your skin or the white part of your eyes** is common with REYATAZ but may be a symptom of a serious problem. These symptoms may be due to increases in bilirubin levels in your blood (bilirubin is made by the liver). Tell your healthcare provider right away if your skin or the white part of your eyes turns yellow.
- **New or worsening diabetes and high blood sugar (hyperglycemia)** have happened in some people who take protease inhibitor medicines like REYATAZ. Some people have had to start taking medicine to treat diabetes or have changes to their dose of their diabetes medicine. Tell your healthcare provider if you notice an increase in thirst or if you start urinating more often while taking REYATAZ.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider if you start having new symptoms after starting REYATAZ.
- **Changes in body fat** can happen in people taking HIV-1 medicines. These changes may include increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the main part of your body (trunk). Loss of fat from the legs, arms, and face may also happen. The exact cause and long-term health effects of these conditions are not known.
- **Increased bleeding problems in people with hemophilia** have happened when taking protease inhibitors like REYATAZ.

The most common side effects of REYATAZ include:

- nausea
- headache
- stomach-area pain
- vomiting
- trouble sleeping
- numbness, tingling, or burning of hands or feet
- dizziness
- muscle pain
- diarrhea
- depression
- fever

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of REYATAZ. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store REYATAZ?

REYATAZ capsules:

- Store REYATAZ capsules at room temperature, between 68°F to 77°F (20°C to 25°C).
- Keep capsules in a tightly closed container.
- The REYATAZ bottle comes with a child-resistant closure.

REYATAZ oral powder:

- Store REYATAZ oral powder at a temperature of 68°F to 86°F (20°C to 30°C).
- Store REYATAZ oral powder in the original packet. Do not open until ready to use.
- After REYATAZ oral powder is mixed with food or liquid it may be kept at a temperature of 68°F to 86°F (20°C to 30°C) for up to 1 hour. Take REYATAZ oral powder within 1 hour after mixing with food or liquid.

Keep REYATAZ and all medicines out of the reach of children.

General information about the safe and effective use of REYATAZ

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use REYATAZ for a condition for which it was not prescribed. Do not give REYATAZ to other people, even if they have the same symptoms that you have. It may harm them. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about REYATAZ that is written for health professionals.

For more information, go to www.reyataz.com or call 1-800-321-1335.

What are the ingredients in REYATAZ?

Active ingredient: atazanavir sulfate

Inactive ingredients:

REYATAZ capsules: crospovidone, lactose monohydrate, and magnesium stearate. The capsule shells contain gelatin, FD&C Blue No. 2, titanium dioxide, black iron oxide, red iron oxide, and yellow iron oxide. The capsules are printed with ink containing shellac, titanium dioxide, FD&C Blue No. 2, isopropyl alcohol, ammonium hydroxide, propylene glycol, n-butyl alcohol, simethicone, and dehydrated alcohol.

REYATAZ oral powder: aspartame, sucrose, and orange-vanilla flavor.

Distributed by:

Bristol-Myers Squibb Company, Princeton, NJ 08543 USA.

REYATAZ® is a registered trademark of Bristol-Myers Squibb Company. Other brands listed are the trademarks of their respective owners and are not trademarks of Bristol-Myers Squibb Company.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: December 2024

[print code]

Instructions for Use
REYATAZ® (RAY-ah-taz)
(atazanavir)
oral powder

Read this Instructions for Use before you prepare your child's first dose of REYATAZ oral powder, each time you get a refill, and as needed. There may be new information. This information does not take the place of talking to your child's healthcare provider about their medical condition or treatment. Ask your child's healthcare provider or pharmacist if you have questions about how to mix or give a dose of REYATAZ oral powder.

Important information:

- For more information about REYATAZ oral powder, see the Patient Information leaflet.
- REYATAZ oral powder must be mixed with food or liquid. If REYATAZ oral powder is mixed with water, your child must eat food right after taking REYATAZ oral powder.
- REYATAZ oral powder must be taken with ritonavir.
- Talk with your child's healthcare provider to help decide the best schedule for giving your child REYATAZ oral powder.

Instructions for mixing REYATAZ oral powder:

REYATAZ oral powder should be mixed with food such as applesauce or yogurt, instead of a liquid (milk, infant formula, or water) in young children and infants who can take food.

- Infants less than 6 months old and who cannot eat solid food or drink from a cup should be given REYATAZ oral powder mixed with infant formula using an oral dosing syringe.
- REYATAZ oral powder that is mixed in infant formula or liquid should not be given using a baby bottle.

When preparing REYATAZ oral powder with either food or liquid, choose a clean, flat work surface. Place a clean paper towel on the work surface. Place the supplies you will need on the paper towel.

Wash and dry your hands before and after preparing REYATAZ oral powder.

Preparing a dose of REYATAZ oral powder mixed with food:

Before you prepare a dose of REYATAZ oral powder mixed with food, gather the following supplies:

- paper towel
- tablespoon
- **small** clean container (such as a **small** cup or bowl)
- a food such as applesauce or yogurt
- the correct number of packets of REYATAZ oral powder needed for the prescribed dose
- a clean pair of scissors

Step 1. Place at least 1 tablespoon of a food such as applesauce or yogurt in the small container (see Figure A).




Figure A

Step 2. Tap the packet of REYATAZ oral powder to settle the contents to the bottom of the packet (see Figure B).




Figure B

Step 3. Using a clean pair of scissors, cut open the packet on the dotted line (see Figure C).




Figure C

Step 4. Empty the contents of the packet into the small container onto the food (see Figure D).




Figure D

Repeat Steps 2 through 4 for each packet of REYATAZ oral powder needed for the total prescribed dose.

Step 5. Use a tablespoon to gently mix the powder and the food together (see Figure E).



Figure E

Steps 6 through 8 must be completed **within 1 hour** of mixing the medicine.

Step 6. Use the tablespoon or a small spoon to feed the REYATAZ oral powder and food mixture to your child. Look in your child's mouth to make sure that all of the mixture is swallowed.

Step 7. Add 1 tablespoon more of food to the empty container and gently stir to mix with any contents that may still be in the container.

Step 8. Use the tablespoon or a small spoon to feed your child the mixture, making sure your child has swallowed all of the mixture.

Step 9. Give your child ritonavir as prescribed right after taking REYATAZ oral powder.

Step 10. Wash the container and tablespoon. Allow the container and spoon to dry. Throw away the paper towel and clean the work surface.

Preparing a dose of REYATAZ oral powder mixed with liquid in a small drinking cup:

Before you prepare a dose of REYATAZ oral powder mixed with liquid in a small drinking cup, gather the following supplies:

- paper towel
- spoon
- 30 milliliter (mL) medicine cup (ask your pharmacist for this). See Figure F.
- small drinking cup
- liquid such as milk or water
- the correct number of packets of REYATAZ oral powder needed for the prescribed dose
- a clean pair of scissors




Figure F

Step 1. Using the 30 mL medicine cup, pour at least 30 mL of liquid into the small drinking cup (see Figure G).



Figure G

Step 2. Tap the packet of REYATAZ oral powder to settle the contents to the bottom of the packet (see Figure H).




Figure H

Step 3. Using a clean pair of scissors, cut open the packet on the dotted line (see Figure I).




Figure I

Step 4. Empty the contents of the packet into the small drinking cup (see Figure J).



Figure J

Repeat Steps 2 through 4 for each packet of REYATAZ oral powder needed for the total prescribed dose.

Step 5. Hold the small drinking cup with one hand. With your other hand, use the spoon to gently mix the powder and the liquid (see Figure K).




Figure K

Steps 6 and 7 must be completed **within 1 hour** of mixing the medicine.

Step 6. Have your child drink all of the mixture in the small drinking cup.

Step 7. To make sure there is no mixture left in the small drinking cup add 15 mL more liquid to the small drinking cup:

- Stir with the spoon.
- **Repeat Step 6 above.**

If REYATAZ oral powder is mixed with water, your child must eat food right after taking REYATAZ oral powder.

Step 8. Give your child ritonavir as prescribed right after taking REYATAZ oral powder.

Step 9. Wash the small drinking cup, medicine cup, and spoon. Allow the small drinking cup, medicine cup, and spoon to dry. Throw away the paper towel and clean the work surface.

Preparing a dose of REYATAZ oral powder mixed with liquid infant formula using an oral dosing syringe and a small medicine cup:

Before you prepare a dose of REYATAZ oral powder mixed with infant formula using an oral dosing syringe, gather the following supplies:

- paper towel
- small spoon
- 30 milliliter (mL) medicine cup (ask your pharmacist for this). See Figure L.
- 10 mL oral dosing syringe (ask your pharmacist for this). See Figure L.
- infant formula
- the correct number of packets of REYATAZ oral powder needed for the prescribed dose
- a clean pair of scissors




Figure L

Step 1. Prepare the infant formula according to the directions on the infant formula package.

Step 2. Pour 10 mL of infant formula into the medicine cup (see Figure M).




Figure M

Step 3. Tap the packet of REYATAZ oral powder to settle the contents to the bottom of the packet (see Figure N).




Figure N

Step 4. Using a clean pair of scissors, cut open the packet on the dotted line (see Figure O).




Figure O

Step 5. Empty the contents of the packet into the medicine cup (see Figure P).




Figure P

Repeat Steps 3 through 5 for each packet of REYATAZ oral powder needed for the total prescribed dose.

Step 6. Hold the medicine cup with one hand. With your other hand, use the small spoon to gently mix the powder and the infant formula (see Figure Q).



Figure Q

Steps 7 through 9 must be completed **within 1 hour** of mixing the medicine.

Step 7. Draw up the powder and infant formula mixture into the oral dosing syringe as follows:

- Check that the plunger is completely pushed into barrel of the syringe (see Figure R).




Figure R

- Place the tip of the syringe into the powder and infant formula mixture in the medicine cup (see Figure S).




Figure S

- Slowly pull back on the plunger and draw up 10 mL of the mixture (see Figure T).




Figure T

Step 8. Place the tip of the oral dosing syringe in your baby's mouth along the inner cheek on either the right or left side (see Figure U). Slowly push on the plunger to give your baby all of the REYATAZ oral powder and infant formula mixture.

- Draw up any remaining mixture with the oral dosing syringe and repeat until all of the mixture has been given to the baby.




Figure U

Step 9. To make sure there is no mixture left in the medicine cup or syringe:

- Repeat Step 1 above to add 10 mL more infant formula to the medicine cup.
- Stir with a small spoon.
- **Then repeat Steps 7 through 8 above.**

To make sure that your baby gets all of the medicine, do not give REYATAZ oral powder in a baby bottle.

Step 10. Give your baby ritonavir as prescribed right after taking REYATAZ oral powder.

Step 11. Remove the plunger from the oral dosing syringe. Wash the medicine cup, spoon, and oral dosing syringe. Allow the medicine cup, spoon, and oral dosing syringe to dry. Throw away the paper towel and clean the work surface.

How should I store REYATAZ oral powder?

- Store REYATAZ oral powder at a temperature of 68°F to 86°F (20°C to 30°C).
- Store REYATAZ oral powder in the original packet. Do not open until ready to use.
- After REYATAZ oral powder is mixed with food or liquid, it may be kept at a temperature of 68°F to 86°F (20°C to 30°C) for up to 1 hour. Take REYATAZ oral powder within 1 hour after mixing with food or liquid.

Keep REYATAZ oral powder and all medicines out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Distributed by:
Bristol-Myers Squibb Company
Princeton, NJ 08543 USA

[print code]

Revised: September 2020